

Technical Information

Nutrient Agar 1.5%

Product Code : DM1087I

Application: Nutrient Agar 1.5% recommended Nutrient Agar 1.5% is a general purpose nutrient medium which can be used for cultivation of fastidious microorganisms after appropriate enrichment.

Composition**

Ingredients	Gms / Litre
Beef extract	3.000
Peptic digest of animal tissue	5.000
Sodium chloride	5.000
Agar	15.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Nutrient Agar is recommended for cultivation and maintenance of nonfastidious microorganisms. Recently ISO Committee with a slight modification⁽²⁾ has recommended this medium (DM1087I) for subcultivation of Pseudomonas species isolated from meat and meat products.

Peptic digest of animal tissue is the principal source of organic nitrogen while Beef extract provides carbohydrates, vitamins, organic nitrogen compounds and salts. Sodium chloride makes the medium isotonic preventing haemolysis of red blood corpuscles. This Nutrient Agar may be used for blood culturing work after the addition of sterile 5% v/v defibrinated blood and Sodium chloride (3g/l)^(1,3).

Methodology

Suspend 28.0 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. If desired, it can be enriched with sterile blood, ascitic fluid or serum after cooling medium base to 45-50°C.

Quality Control

Physical Appearance

Cream to yellow coloured homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Yellow coloured clear gel forms in Petri plates. With the addition of blood, cherry red coloured, opaque gel forms in petri plates.

Reaction

Reaction of 2.8% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH Range 6.80-7.20

Cultural Response/ characteristics

DM 1087I: Cultural characteristics observed after an incubation at 35-37°C for 18 - 24 hours.

Organism	Inoculum (CFU)	Growth	Recovery
Enterococcus faecalis ATCC 29212	50-100	luxuriant	>=70%
Escherichia coli ATCC 25922	50-100	luxuriant	>=70%
Pseudomonas aeruginosa ATCC 27853	50-100	luxuriant	>=70%
Staphylococcus aureus ATCC 25923	50-100	luxuriant	>=70%
Streptococcus pyogenes ATCC 19615	50-100	luxuriant	>=70%
Streptococcus pneumoniae ATCC 6303	50-100	luxuriant	>=70%



Dehydrated Culture Media
Bases / Media Supplements

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Speck M. (Ed.), 1984, Compendium of Methods for the Microbiological Examination of Foods, 2nd ed., APHA, Washington D.C.
2. International Organization for Standardization (ISO), 1995, Draft ISO/DIS 13720.
3. Pelczar, Chan and Kreig, 1986, Microbiology, 5th ed., McGraw Hill Book Co., N.Y.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

