



Dehydrated Culture Media
Bases / Media Supplements

Technical Information

Sabouraud Maltose Broth

Product Code: DM 1064

Application: Sabouraud Maltose Broth is used as an excellent medium for the propagation of moulds and yeasts, particularly the parasitic fungi concerned with skin and scalp lesions.

Composition**

Ingredients	Gms / Litre
Mycological, peptone	10.000
Maltose	40.000
Final pH (at 25°C)	5.6±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Fungi were among the first microorganisms recognized without a microscope because some of the fruiting structures, such as the mushrooms, are large enough to be seen. Fungi can be grouped simply on the basis of morphology as either yeasts or moulds ⁽¹⁾. Sabouraud Maltose Broth was formulated by Sabouraud ⁽²⁾ and is used for the isolation and differentiation of yeast and moulds ⁽³⁻⁵⁾.

Mycological peptone provides nitrogen, vitamins, minerals, amino acids and growth factors. Maltose provides an energy source for the growth of microorganisms. The low pH favours fungal growth and inhibits contaminating bacteria from clinical specimens ⁽¹⁾. For isolation of fungi from contaminated specimens, a selective medium should be inoculated simultaneously. Incubate cultures for 4 to 6 weeks before reporting as negative.

Methodology

Suspend 50 grams of powder media in 1000 ml distilled water. Shake well & heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and dispense into sterile test tubes.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light amber coloured clear solution in tubes

Reaction

Reaction of 5.0% w/v aqueous solution at 25°C. pH : 5.6±0.2

pH Range:- 5.40-5.80

Cultural Response/Characteristics

DM1064: Cultural characteristics observed after an incubation at 25 - 30°C for 48-72 hours .(Incubate Trichophyton species for upto 7 days)

Organism	Inoculum (CFU)	Growth
<i>Aspergillus brasiliensis</i> ATCC 16404	50-100	good-luxuriant
<i>Candida albicans</i> ATCC 10231	50-100	good-luxuriant
<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant
<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	good-luxuriant
<i>Trichophyton rubrum</i> ATCC 28191	50-100	good-luxuriant
<i>Lactobacillus casei</i> ATCC 9595	50-100	good-luxuriant



Storage and Shelf Life-

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Murray P. R., Baron J. H., Pfaller M. A., Tenover J. C. and Tenover F. C., (Ed.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
2. Sabouraud R., 1892, Ann. Dermatol. Syphil. 3 : 1061.
3. Davidson and Dowding, 1932, Arch. Dermatol. Syphilol. 26:660.
4. Davidson, Dowding and Buller. 1932. Can. J. Res. 6:1.
5. Frank L. S., 1932, Arch. Dermatol. Syphilol., 26: 457

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.