



Dehydrated Culture Media  
Bases / Media Supplements

## Technical Information

### Tryptose

**Product Code: BA2030**

#### Principle & Interpretation

Tryptose is specially prepared enzymic hydrolysate of protein, to meet the nutritional requirements of fastidious microorganisms. It is used in Tryptose media, where blood cells are maintained in excellent state and so haemolytic reactions can be easily demonstrated. It is also used in vaccine preparation media for rapid and luxuriant growth as desired for large scale production of vaccines and toxins, for supplementing tissue culture media .

#### Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Safety guidelines may be referred in individual safety data sheets.

#### Limitations

1. It is biological origin product since variation in colour of powder and clarity may observed.
2. Each lot of the product has been tested for the organisms specified on the COA. It is recommended to users to validate the medium for any specific microorganism other than mentioned in the COA based on the user 's requirement.
3. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium prepared by the product.

#### Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature

#### Quality Control

##### Appearance

Light yellow to yellow homogeneous free flowing powder characteristic but not putrescent

##### Solubility

Freely soluble in distilled/purified water, insoluble in alcohol.

##### Clarity

1% w/v aqueous solution is clear without any haziness after autoclaving at 15 lbs pressure (121°C) for 15 minutes.

##### pH

pH of 2% w/v aqueous solution at 25°C 6.2 - 7.2

##### Microbial Load :

Bacterial Count : <= 2000 CFU/gram by plate method, when incubated at 30-35°C for not less than 3 days

Yeast & mould Count : <= 100 CFU/gram by plate method, when incubated at 20-25°C for not less than 5 days.

##### Test for pathogens :

1. *Escherichia coli*- Absent/gram of sample
2. *Salmonella* species- Absent/10 gram of sample



Dehydrated Culture Media  
Bases / Media Supplements

3. *Pseudomonas aeruginosa*- Absent/gram of sample

4. *Staphylococcus aureus*- Absent/gram of sample

5. *Candida albicans*- Absent/gram of sample

6. *Clostridia*- Absent/gram of sample

**Indole Test** : Tryptophan content: Passes

**Cultural response** : Cultural response observed after an incubation at 35-37°C for 18-24 hours by preparing Lauryl Tryptose Broth (DM1080) using Tryptose as an ingredient.

#### Cultural response

Organism	Growth	Gas Production
<i>Escherichia coli</i> ATCC 25922 (WDCM 00013)	Luxuriant	Positive reaction
* <i>Klebsiella aerogenes</i> ATCC 13048(WDCM 00175)	Luxuriant	Positive reaction
<i>Enterococcus faecalis</i> ATCC 29212 (WDCM00087)	Inhibited	--
<i>Salmonella enterica</i> subsp. <i>Enterica</i> Typhimurium ATCC 14028 (WDCM 00031)	Luxuriant	Negative reaction
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC25923 (WDCM 00034)	Inhibited	--

(\* ) formerly known as *Enterococcus aerogenes*

#### Chemical Analysis

Total Nitrogen	: ≤ 11.80 %
Amino Nitrogen	: ≤ 4.00 %
Sodium chloride	: ≤ 5.00 %
Loss on drying	: ≤ 5.00 %
Residue on ignition	: ≤ 15.00 %

## Storage and Shelf Life

Store between 10-30°C in tightly closed container and away from bright light. Use before expiry date on label. On opening, product should be properly stored in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use

## Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.