

Technical Information

MacConkey Agar Base

Product Code: DM 2024

Application: - MacConkey Agar Base is recommended for studying fermentation reactions of coliforms by adding carbohydrate.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	17.000
Proteose peptone	3.000
Bile salts	1.500
Sodium chloride	5.000
Neutral red	0.030
Crystal violet	0.001
Agar	13.500
Final pH (25°C)	7.1±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

MacConkey Agar is one of the earliest selective and differential medium for cultivation of enteric microorganisms from large number of clinical specimens ^(1, 2). MacConkey Agar Base is used for studying carbohydrate fermentation reactions of coliforms by adding carbohydrates either individually or in combination ⁽³⁾.

MacConkey Agar Base has peptic digest of animal tissue and proteose peptone, which provide nitrogen, carbon and vitamin source for the growth of bacteria. This medium does not contain carbohydrates. However for studying fermentation reaction, carbohydrate of interest has to be added while preparing medium. The selective action of this medium is due to bile salts and crystal violet, which are inhibitory to most of the species of gram-positive bacteria. Gram-negative bacteria usually grow well on the medium and are differentiated by their ability to ferment carbohydrates. Carbohydrate (lactose) fermenting strains grow as red or pink and may be surrounded by a zone of acid precipitated bile. The red colour is due to production of acid from carbohydrate, absorption of neutral red and subsequent colour change of the dye when the pH of the medium falls below 6.8. Sodium chloride helps to maintain osmotic balance.

Methodology

Suspend 40 grams of powder media in 1000 ml distilled water. Add 10 grams of lactose or other carbohydrates of choice. Shake well & heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure for 15 minutes. Mix well and pour into sterile Petri plates.

Quality Control

Physical Appearance

Light yellow to pink homogeneous free flowing powder

Gelling

Firm, comparable with 1.35% Agar gel

Colour and Clarity of prepared medium

Red with purplish tinge clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 4.0% w/v aqueous solution at 25°C, pH: -7.1±0.2

pH range 6.90-7.30

Cultural Response/ characteristics

DM 2024: Cultural characteristics observed with added 1% lactose, after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
Escherichia coli ATCC 25922	50-100	luxuriant	≥50 %	pink to red with bile precipitate pink to red
Enterobacter aerogenes ATCC 13048	50-100	luxuriant	≥50 %	pink to red
Enterococcus faecalis ATCC 29212	50-100	fair to good	30-40%	pale pink to red
Proteus vulgaris ATCC 13315	50-100	luxuriant	≥50 %	colourless
Salmonella Paratyphi A ATCC 9150	50-100	luxuriant	≥50 %	colourless
Shigella dysenteriae ATCC 13313	50-100	fair to good	30-40%	colourless
Salmonella Paratyphi B ATCC 8759	50-100	luxuriant	≥50 %	colourless
Salmonella Enteritidis ATCC 13076	50-100	luxuriant	≥50 %	colourless
Salmonella Typhi ATCC 6539	50-100	luxuriant	≥50 %	colourless
Staphylococcus aureus ATCC 25923	≥10 ³	inhibited	0%	

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. MacConkey, 1900, The Lancet, ii:20.
2. MacConkey, 1905, J. Hyg., 5:333.
3. Holt, Harris and Teague, 1916, J. Infect. Dis., 18:596.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

Replace date 29-Nov-2025