

Technical Information

Yeast Malt Agar (YM Agar) (ISP Medium No. 2)

Product Code: DM 1424

Application: Yeast Malt Agar (YM Agar) is used for the isolation and cultivation of yeasts, moulds and other aciduric microorganisms.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	5.000
Yeast extract	3.000
Malt extract	3.000
Dextrose	10.000
Agar	20.000
Final pH (at 25°C)	6.2±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Wickerham formulated Yeast Malt Agar for isolation and cultivation of yeasts, moulds and other aciduric microorganisms^(1, 2) Fungistatic materials such as sodium propionate and diphenyl are added to YM Agar to eliminate moulds and permit the enumeration of yeasts from mixed population. YM Agar is also recommended by APHA⁽³⁾.

Wickerham suggested the use of Yeast Malt Broth (DM1425) as an enrichment medium for yeasts by adding a layer of sterile paraffin oil (about 1 cm) on the surface of inoculated broth. After the growth occurs it should be streaked on YM Agar to obtain isolated colonies of fermentative species. To isolate fermentative as well as oxidative strains, acidified YM Broth (DM1425) is placed on a rotary shaker for 1 or 2 days which favors development of yeast cells while the sporulation of moulds is prevented and yeasts can be isolated by streaking on YM Agar. Peptic digest of animal tissue serves as a source of carbon, nitrogen and essential nutrients. Yeast extracts supplies vitamin B complex nutrients and other growth factors. Malt extract serves as an additional source of carbon. Dextrose is the carbohydrate and energy source. To increase the selectivity, the media can be acidified by the addition of sterile 10% HCl, tartaric acid or 10% citric acid. Alternatively antibiotics (penicillin 20U/ml or streptomycin to a final concentration of 40mcg/ml) can be added. Acidified agar medium should not be reheated.

Methodology

Suspend 20.5 grams of powder media in 490 ml distilled water. Shake well & heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. For preparing selective media acidify the media up to pH 3.0 to 4.0 by aseptically adding 1 vial of 10% Lactic Acid Solution (MS2095). DO NOT HEAT the media after addition of acid. Mix well and pour into sterile Petri plates.

Quality Control

Physical Appearance

Cream to beige homogeneous free flowing powder

Gelling

Firm, comparable with 2.0% Agar gel.

Colour and Clarity of prepared medium

Light amber coloured clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 4.1% w/v aqueous solution at 25°C. pH : 6.2±0.2

pH Range:-

6.00-6.40

Cultural Response/Characteristics



Dehydrated Culture Media
Bases / Media Supplements

DM 1424: Cultural characteristics observed after an incubation at 25-30°C for 40-72 hours.

Organism	Inoculum	Growth at pH 3.4	Growth at pH 6.2	Recovery
Cultural Response				
*Aspergillus brasiliensis ATCC 16404	50-100	Good-luxuriant	Good-luxuriant	>=50%
Candida albicans ATCC 10231	50-100	Good-luxuriant	Good-luxuriant	>=50%
Escherichia coli ATCC 25922	50-100	Inhibited	Good-luxuriant	>=50%
Lactobacillus casei ATCC 9595	50-100	Poor	Good-luxuriant	>=50%
Lactobacillus leichmannii ATCC 4797	50-100	Poor	Good-luxuriant	>=50%
Saccharomyces cerevisiae ATCC 9763	50-100	Good-luxuriant	Good-luxuriant	>=50%

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Wickerham L. J., 1951, U.S. Dept. Agric. Tech. Bull. No.1029.
2. Wickerham L. J., 1939, J. Tropical Med. Hyg., 42:176.
3. Downes F. P. and Ito K.,(Ed.), 2001, Compendium of Methods for the Microbiological examination of Foods, 4th Ed, APHA Inc. Washington DC.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.

Replace Date 3 08-Mar-2019

