

### **Technical Information**

# Luria Bertani Agar Plate w/100µg/ml Ampicillin

### Product Code: PM 2151A

Application: Recommended for the cultivation and maintenance of recombinant strains of E.coli for genetic and molecular biology studies

#### Composition\*\*

| Ingredients                                                     | Gms / Litre |
|-----------------------------------------------------------------|-------------|
| Tryptone                                                        | 10.000      |
| Yeast extract                                                   | 5.000       |
| Sodium chloride                                                 | 10.000      |
| Agar                                                            | 15.000      |
| Ampicillin                                                      | 100mg       |
| Final pH ( at 25°C)                                             | 7.5±0.2     |
| **Formula adjusted, standardized to suit performance parameters |             |

### **Principle & Interpretation**

Luria Bertani Agar is prepared as described by Lennox (4) for cultivation and maintenance of recombinant strains of Escherichia coli. Luria Bertani Agar, Miller (1) is slightly different with double amount of sodium chloride. The media is nutritionally rich for the growth of pure cultures of recombinant strains. Strains derived from Escherichia coli K12 are deficient in Vitamin B synthesis are further modified by specific mutation to create auxotrophic strains and are therefore unable to grow on nutritionally deficient media.

Tryptone provides peptides and peptones while Vitamin B complex is provided by yeast extract. Sodium chloride provides sodium ions for membrane transport and also maintains the osmotic equilibrium of the medium.

# Type of specimen

Recombinant strains of E.coli

# Specimen Collection and Handling

For Recombinant strain samples follow appropriate techniques for handling specimens as per established guidelines (1,4). After use, contaminated materials must be sterilized by autoclaving before discarding.

# Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

#### Limitations

- 1. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium
- 2. Each lot of the medium has been tested for the organisms specified on the COA. It is recommended to users to validate the medium for any specific microorganism other than mentioned in the COA based on the user's unique requirement.
- 3. Further biochemical and serological tests must be carried out for complete identification.



### Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

### Methodology

Either streak, inoculate or surface spread the test inoculum (50-100 CFU) aseptically on the plate.

### **Quality Control**

#### Appearance

Sterile Luria Bertani Agar Plate w/100µg/ml Ampicillin in 90mm disposable plates.

Reaction

7.30-7.70

Sterility test

Passes release criteria

**Cultural Response** 

Cultural characteristics observed after an incubation at 35-37°C for 18 - 24 hours.

Oragnism Growth

Escherichia coli ATCC23724 Luxuriant
Escherichia coli ATCC25922 (00013\*) Luxuriant
Escherichia coli DH5 alphaMTCC 1652 Luxuriant

Key: \* - Corresponding WDCM numbers

# Storage and Shelf Life

On receipt store between 2-8°C Use before expiry date on the label. Product performance is best if used within stated expiry period.

# Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (2,3)

# **Further Reading**

- 1. Atlas R.M., 1993, Handbook of Microbiological Media, Ed. by Parks L., CRC Press, Inc.
- $2. \ \ \text{Isenberg, H.D. Clinical Microbiology Procedures Handbook } 2^{\mbox{nd}} \ \ \mbox{Edition}.$
- 3. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
- 4. Lennox E.S., 1955, Transduction of Linked Genetic Characters of the host by bacteriophage P1., Virology, 1:190.



### **Disclaimer**

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing
- of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performens parameters.