



Ready Prepared Media

Technical Information

Sabouraud Chloramphenicol Agar Plate w/ 1% Glycerol (γ irradiated) (Triple Pack)

Product Code: PM 2946GT

Application: Recommended for the selective cultivation of yeasts and moulds

Composition**

Ingredients	Gms / Litre
Tryptone	5.000
Peptone	5.000
Dextrose (Glucose)	40.000
Chloramphenicol	0.050
Agar	15.000
Glycerol	10 ml
Final pH (at 25°C)	5.6±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Sabouraud Chloramphenicol Agar is cited as Medium C and recommended for cultivation of yeasts and moulds. This medium was described originally by Sabouraud (7) for the cultivation of fungi, particularly useful for the fungi associated with skin infections. The medium is often used with antibiotics such as Chloramphenicol (1) for the isolation of pathogenic fungi from materials containing large numbers of fungi or bacteria.

Tryptone and peptone provide nitrogenous and carbonaceous compounds, long chain amino acids, and other essential growth nutrients. Dextrose provides an energy source. Chloramphenicol inhibits a wide range of Gram-positive and Gram-negative bacteria which makes the medium selective for fungi (5). The low pH favors fungal growth and inhibits contaminating bacteria from clinical specimens (6).

Type of specimen

Food and dairy samples

Specimen Collection and Handling

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (2,8,9).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

Read the label before opening the pack. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.



Ready Prepared Media

Limitations

1. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium .
2. Each lot of the medium has been tested for the organisms specified on the COA. It is recommended to users to validate the medium for any specific microorganism other than mentioned in the COA based on the user's unique requirement.
3. Presence of chloramphenicol may inhibit certain pathogenic fungi.
4. It is recommended to store the plates ta 24-30°C to avoid minimum condensation.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Methodology

Either streak, inoculate or surface spread the test inoculum (50-100 CFU) aseptically on the plate. Caution: Some pathogenic fungi may produce infective spores which are easily dispersed in air, so examination should be carried out in safety cabinet.

Quality Control

Appearance

Sterile Sabouraud Chloramphenicol Agar Plate w/ 1% Glycerol in 90 mm disposable plates.

Colour

Light amber coloured medium

Quantity

30 ml of medium in 90mm disposable plate

pH

5.40-5.80

Dose of irradiation (Kgy)

13.00- 20.00

Sterility Check

Passes release criteria

Cultural Response

Cultural characteristics observed after an incubation at 20-25°C for 48-72 hours (Incubate for 7 days for *Trichophyton* species).

Organism	Inoculum(CFU)	Growth	Recovery
<i>Aspergillus brasiliensis</i> ATCC 16404 (00053*)	50-100	Good luxuriant	
<i>Candida albicans</i> ATCC 10231 (00054*)	50-100	Good luxuriant	>=50%
<i>Escherichia coli</i> ATCC 25922 (00013*)	>=10 ⁴	Inhibited	0%
<i>Lactobacillus casei</i> ATCC 334	>=10 ⁴	Inhibited	0%
<i>Saccharomyces cerevisiae</i> ATCC 9763 (00058*)	50-100	Good luxuriant	>=50%
<i>Trichophyton rubrum</i> ATCC28191	50-100	Good luxuriant	
<i>Escherichia coli</i> NCTC 9002	>=10 ⁴	Inhibited	0%
<i>Escherichia coli</i> ATCC 8739 (00012*)	>=10 ⁴	Inhibited	0%

Key : (*) - Corresponding WDCM numbers



Ready Prepared Media

Storage and Shelf Life

- On receipt store between 20-30°C Use before expiry date on the label.
- Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (3,4).

Further Reading

1. Ajello L., 1957, J. Chron. Dis., 5:545.
2. American Public Health Association, Standard Methods for the Examination of Dairy Products, 1978, 14th Ed., Washington D.C.
3. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
4. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
5. Lorian (Ed.), 1980, Antibiotics In Laboratory Medicine, Williams and Wilkins, Baltimore.
6. Murray, P. R 2005, In Manual of Clinical Microbiology, 7th ed., ASM, Washington, D.C.
7. Sabouraud K., 1892, Ann. Dermatol. Syphilol, 3:1061.
8. Salfinger Y., and Tortorello M.L., 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
9. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.