

Technical Information

Linsmaier & Skoog Vitamins (100X)

Product Code: VP1017

Application: Linsmaier & Skoog vitamins (100X) powder is ready to use mixture of vitamins which is sufficient to prepare 100 litres of medium from content of one vial. The vial contains 10.04 gm of powdered vitamin mixture.

Composition**

Ingredients	mg/Litre
myo - Inositol	100.00
Thiamine hydrochloride	0.40
TOTAL gm/litre	0.10

Methodology

Suspend 0.10 grams of dehydrated vitamins powder# in 600ml of distilled water. Apply constant gentle stirring to the solution till the powder dissolves completely. Add desired heat stable supplements prior to autoclaving. Adjust the medium to the desired pH using 1N HCl/NaOH. Make up the final volume to 1000ml with distilled water. Sterilize the medium by autoclaving at 15 lbs or 121°C for 15 minutes. Cool the autoclaved medium to 45°C before adding the filter sterilized heat labile supplements. Dispense the desired amount of medium aseptically in sterile culture vessels

Weight after vacuum drying to remove all water

Quality Control

Appearance

White to off-white, homogeneous, free flowing powder.

Solubility

10.04gm/litre soluble in distilled water.

Colour and clarity

Colourless to light yellow, clear solution

pH at 25°C:

3.9 ±0.5 of 1.004% w/v dehydrated powdered vitamins mixture.

Cultural Response

Cultural condition :

Incubation period : 5 weeks
 Relative humidity : 60% ± 2%
 Temperature : 22°C ± 2°C
 Photoperiod (D:N) in hours : 16:8

Cell Line	Type of Culture	Results
<i>Musa</i> species	Shoot culture	No structural deformity observed No necrotic tissues, Actively growing shoots, No toxicity to shoots
<i>Daucus</i> species	Callus culture	No necrotic tissues, Actively growing callus, No toxicity to callus

[The medium is prepared as per direction. The growth promoting activity of this vitamin mixture is evaluated using two plant species viz. *Musa* species and *Daucus* species through three passages. Plant growth hormones (e.g. 2,4-D, NAA, Kinetin and 6-BAP) are added in suitable combinations and concentrations.]

Storage and Shelf Life

Powdered vitamin mixtures are very hygroscopic and must be protected from atmospheric moisture. It is advisable to use the entire contents of vial immediately after opening. In case of unused powder should be stored in desiccator and refrigerated at 2-8°C.

Use before the expiry date.

Reference

- Linsmaier E.M. & Skoog F., *Physiol. Plant.*, (1965), 18, 100 - 127

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.