

Technical Information

De Greef & Jacobs Plant Salt Mixture

Product Code: TS2114

Composition**

Ingredients	mg/Litre
Potassium nitrate	2000.00
Ammonium sulphate	400.00
Calcium chloride.2H ₂ O	300.00
Magnesium sulphate	244.17
Sodium phosphate monobasic	250.00
Potassium chloride	600.00
Manganese sulphate.H ₂ O	1.68
Boric acid	10.62
Potassium iodide	1.58
Molybdic acid (sodium salt).2H ₂ O	0.0025
Zinc sulphate.7H ₂ O	1.06
Copper sulphate.5H ₂ O	0.0025
Cobalt chloride.6H ₂ O	0.0025
Ferrous sulphate.7H ₂ O	27.80
EDTA disodium salt.2 H ₂ O	37.26
TOTAL	3.87 gm/litre

Principle And Interpretation

De Greef & Jacobs plant salt mixture has been specially formulated for plant cell, tissue and organ cultures. The mixture contains macroelements, microelements and iron source.

Directions

Suspend 3.78 grams of dehydrated plant salt mixture# in 600ml of distilled water and rinse media vial with small quantity of distilled water to remove traces of powder. Apply constant gentle stirring to the solution till the powder dissolves completely. Add desired heat stable supplements prior to autoclaving. Adjust the medium to the desired pH using 1N HCl/NaOH. Make up the final volume to 1000ml with distilled water. Sterilize the medium by autoclaving at 15 lbs or 121°C for 15 minutes. Cool the autoclaved medium to 45°C before adding the filter sterilized heat labile supplements. Dispense the desired amount of medium aseptically in sterile culture vessels.

Weight after vacuum drying to remove all water

Quality Control

Appearance	: White to off-white, homogeneous, free flowing powder.
Solubility	: 3.78 gm/litre soluble in distilled water.
Colour and Clarity	: Colourless to light yellow, clear solution.
pH at 25°C	: 4.0 ± 0.5 (under observations) of 0.378% w/v dehydrated plant salt mixture.

Cultural Response :

Cultural condition :

- Incubation period : 5 weeks
- Relative humidity : 60% ± 2%
- Temperature : 22°C ± 2°C
- Photoperiod (D:N) in hours : 16:8

Cell Line	Types Of Culture	Results
<i>Musa species</i>	Shoot culture	No structural deformity observed No necrotic tissues, Actively growing shoots, No toxicity to shoots
<i>Daucus species</i>	Callus culture	No necrotic tissues, Actively growing callus, No toxicity to callus

[The medium is prepared as per direction. The growth promoting activity of this plant salt mixture is evaluated using two plant species viz. *Musa species* and *Daucus species* through three passages. Plant growth hormones (e.g. 2,4-D, NAA, Kinetin and 6-BAP) are added in suitable combinations and concentrations.]

Storage and Shelf Life

Dehydrated plant salt mixture powder is extremely hygroscopic and should be protected from atmospheric moisture. If possible, the entire content of each bottle should be used immediately after opening or else the unused portion should be stored in a desiccator and refrigerated at 28°C. Use before the expiry date.

Further Reading

- De Greef W. & Jacobs M., Plant Sci. Lett., (1979), 17, 55 - 61

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performs parameters.