

Technical Information

Stag Horn Fern Multiplication Medium With Vitamins, Sucrose, Adenine sulphate, IAA and Agar

Product Code: PT1119

Application: Stag Horn Fern Multiplication Medium has been developed on the basis of Murashige and Skoog medium for the *in vitro* propagation of fern. Stag horn fern belongs to *Platycerium* genus, family *Polypodiaceae*. The formulation is a nutrient blend of inorganic salts, vitamins, carbohydrate, plant growth regulators and gelling agent.

Stag Horn Fern Multiplication Medium provides all the essential macroelements and microelements. Potassium nitrate and ammonium nitrate serve as sources of nitrogen that promotes morphogenesis. This mixture of cation and anion also maintains pH of medium. Potassium dihydrogen phosphate along with sodium dihydrogen phosphate serve as sources of phosphate. Microelements like Boron, Manganese, Molybdenum, Copper, Iron and Zinc play vital role in the plant metabolism. Thiamine, pyridoxine, nicotinic acid and inositol act as enzymatic cofactors in the universal pathways including glycolysis and TCA cycle along with primary and secondary metabolism in the plants. Adenine sulphate promotes shoot proliferation and multiplication while IAA induces cell division and rooting.

The product is plant tissue culture tested but it is the sole responsibility of the user to ensure the suitability of the medium for individual species.

Composition**

Ingredients	mg/Litre
MACROELEMENTS	
Ammonium nitrate	1650.000
Calcium chloride dihydrate	332.200
Magnesium sulphate anhydrous	180.690
Potassium dihydrogen phosphate	170.000
Potassium nitrate	1900.000
MICROELEMENTS	
Boric acid	6.200
Cobalt chloride hexahydrate	0.025
Copper sulphate pentahydrate	0.025
EDTA disodium salt dihydrate	37.300
Ferrous sulphate heptahydrate	27.800
Manganese sulphate monohydrate	16.897
Potassium Iodide	0.830
Sodium dihydrogen phosphate	147.870
Sodium molybdate	0.213
Zinc sulphate heptahydrate	8.600
VITAMINS	
myo-Inositol	100.000
Nicotinic acid	1000.000
Pyridoxine hydrochloride	1000.000
Thiamine hydrochloride	0.400
CARBOHYDRATE	
Sucrose	30000.000
GELLING AGENT	
Agar	8000.00
OTHERS	
Adenine sulphate	80.000
Indole-3-acetic acid (IAA)	15.000
TOTAL	42.7 ms/litre

Material required but not provided

- Autoclaved distilled water
- 1N NaOH/HCl

Quality Control

Appearance

White to off-white, homogenous, free flowing powder

Solubility

42.7 gms/litre soluble after boiling in distilled water

Colour and Clarity

Colourless to light yellow solution, hazy gel is formed on cooling

pH at 25°C

3.50-4.50

Plant Tissue Culture Test

The growth promoting properties of medium is assessed by providing plant cultures with relative humidity of about 60%±2%, temperature 22°C±2°C and photoperiod of about 16:8. The plant species showed actively growing callus and shoots with no structural, necrotic and toxic deformity.

Directions

- Reconstitute medium by adding required quantity of powder in two-third of total volume with constant, gentle stirring till the medium gets completely dissolved.
- Add heat stable supplements prior to autoclaving.
- Make up the final volume with distilled water.
- Adjust the pH of the medium to 5.75 ± 0.5 using 1N NaOH/HCl.
- Heat the medium to boiling till complete dissolution of gelling agent.
- Sterilize the medium by autoclaving at 15 lbs and 121°C for 15 min.
- Cool the autoclaved medium to about 45°C before adding heat labile supplements.
- Aseptically dispense the desired amount of medium under a laminar airflow unit in sterile culture vessels

Storage and Shelf Life

- The plant tissue culture medium powder is extremely hygroscopic and must be stored at 2-8°C in air tight containers.
- Preferably, entire content of each package should be used immediately after opening.
- Use before the expiry date.

Precautions

- Ensure appropriate pH of the medium before addition of gelling agent as acidic pH will lead to decreased gelation resulting in semi solid flowing gel while alkaline pH will lead to formation of hardened gel.
- Use of Distilled water/Tissue culture grade water is recommended for media preparation as tap water or lower grade water may lead to salt precipitation and improper gelation.
- Avoid preparation of concentrated solutions, as it will lead to precipitation of salts.

Disclaimer

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.