

## Technical Information

### Phyto Pst Agar Base

**Product Code: PHM1011**

**Application:** Semi-selective medium for the detection of *Pseudomonas syringae pv.tomato* on seeds of tomato.

### Composition\*\*

Ingredients	Grams/Litre
Proteose peptone	20.00
Di-potassium hydrogen phosphate	1.50
Magnesium sulphate anhydrous	0.73
Agar	15.00
Total	37.23 gm/liter

\*\*Formula adjusted standard to suit the performance parameter

### Principle And Interpretation

Bacterial speck of tomatoes is caused by *Pseudomonas syringae pv. tomato*. This bacterium is widespread and is often isolated from plant roots and soil particles since this bacterium is seedborne.. It has been associated with different plants near tomato fields, where it survives as a saprophyte. Recently it was shown that these bacteria may multiply at the base of leaf hairs on healthy tomato leaves and later cause disease. The disease is most serious in cold with high relative humidity.

The foliar symptoms of speck consist of small black lesions, often with a discrete yellow halo. Speck seems to curl the leaves more severely than spot. Bacterial speck lesions are slightly raised, but are generally much smaller. Bacterial speck lesions are very superficial and do not crack or become scaly as in bacterial spot (2).

Proteose peptone provide nitrogenous compounds, carbon, vitamin B complex and trace ingredients. The inorganic phosphates in the medium serve as buffers. The media becomes selective with the addition of CCTP Supplement (PHS1010).

### Directions

Suspend 37.23 grams in 960 ml distilled water. Add 30 ml of 50% glycerol .Dissolve 1.5 g boric acid in 10 ml distilled water. Sterilize the solutions separately by autoclaving at 15 lbs pressure(121°C) for 15 minutes .Cool to 45-50 °C and aseptically add the rehydrated contents of one vial of CCTP supplement (PHS1010) Mix well and pour into Petri plate .

### Quality Control

**Appearance:**

Light yellow coloured, homogeneous, free flowing powder.

**Gelling:**

Firm, comparable with 1.5% Agar gel.

**Colour and Clarity of prepared medium:**

Yellow coloured, opalescent gel forms in Petri plates

**Reaction:**

Reaction of 3.72% w/v aqueous solution is pH 7.5 at 25<sup>o</sup>C

**Cultural Response:**

Cultural characteristics observed after an incubation at 30-32°C for 5-6 days with added CCTP supplement (PHS1010)

Organism (ATCC)	Growth	Colour of the colony
<i>Pseudomonas syringae pv.syringae</i>	luxuriant	small, flat, pink colored
<i>Escherichia coli (25922)</i>	inhibited	-
<i>Staphylococcus aureus (25923)</i>	inhibited	-

### Storage and Shelf Life

Store below 30°C and the prepared medium at 2 - 8°C. Use before expiry date on the label.

### Further Reading

- Ken Pernezny, professor, Plant Pathology Department, Everglades Research and Education Center, Belle Glade, FL; Shouan Zhang, assistant professor, Plant Pathology Department, Tropical Research and Education Center, Homestead; Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611.
- Zitter T. A. Bacterial Diseases of Tomato Vegetable crops.

### Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performs parameters.