Plant Tissue Culture



Product Specification

cdhfinechemical.com

Technical Information

Orchid Maintenance Medium With Calcium Chloride, Vitamins, Sucrose, Tryptone, MES, Activated Charcoal Without Agar

Product Code: PT1055

Application: Orchid Maintenance Medium has been formulated for the effective maintenance of the orchid species.

It is a nutrient blend of inorganic salts, vitamins and carbohydrate. In addition, it is supplemented with MES buffer which maintains optimum buffering and prevents acidification in the media required for the cultivation of orchids. Tryptone is added for the multiplication of protocorm like bodies and differentiation of shoots. Microelements like Boron, Manganese, Molybdenum, Copper, Iron and Zinc enhance metabolism in plants. Boron plays a key role in carbohydrate metabolism. Thiamine, pyridoxine, nicotinic acid act as enzymatic cofactors in universal pathways including glycolysis and TCA cycle along with the primary and secondary metabolism in plants. Activated charcoal adsorbs the inhibitory leachouts from medium.

The product is plant tissue culture tested but it is the sole responsibility of the user to ensure the suitability of the medium for individual species.

Ingredients	mg/Litre
MACROELEMENTS	-
Ammonium nitrate	825.000
Calcium chloride	166.100
Magnesium sulphate	90.340
Potassium nitrate	950.000
Potassium phosphate monobasic	85.000
MICROELEMENTS	
Boric acid	3.100
Cobalt chloride hexahydrate	0.012
Copper sulphate pentahydrate	0.012
EDTA disodium salt dihydrate	37.300
Ferrous sulphate heptahydrate	27.800
Manganese sulphate monohydrate	8.450
Molybdic acid (sodium salt)	0.125
Potassium iodide	0.420
Zinc sulphate heptahydrate	5.300
VITAMINS	
myo-Inositol	100.00

Plant Tissue Culture



Product Specification

cdhfinechemical.com

Nicotinic acid (free acid)	1.000
Pyridoxine HCl	1.000
Thiamine hydrochloride	10.000
CARBOHYDRATE	
Sucrose	20000.000
OTHERS	
Activated charcoal	2000.000
MES	1000.000
Tryptone	2000.000
Total(gms/litre)	27.3

Material required but not provided

- Autoclaved distilled water
- Plant growth regulators
- 1N NaOH/HCl
- Gelling agents like Agar (PCT1901) or CleriGel (PCT1903)

Quality Control

Appearance

Grey to black, homogenous, free flowing powder **Solubility**

27.3 gms/litre soluble in distilled water

Colour and Clarity

Grey to black opaque solution

pH at 25ºC

4.60 - 5.60

Plant Tissue Culture Test

The growth promoting properties of medium is assessed by providing plant cultures with relative humidity of about 60%±2%, temperature 22ºC±2ºC and photoperiod of about 16:8. The plant species showed actively growing callus and shoots with no structural, necrotic and toxic deformity.

Directions

- Reconstitute medium by adding required quantity of powder in two-third of total volume with constant, gentle stirring till the medium gets completely dissolved.
- Add heat stable supplements prior to autoclaving.
- Make up the final volume with distilled water.
- Adjust the pH of the medium to 5.75 ± 0.5 using 1N NaOH/HCl.
- Add gelling agent and heat the medium to boiling till complete dissolution of gelling agent.
- Sterilize the medium by autoclaving at 15 lbs and 121°Cfor 15 min.
- Cool the autoclaved medium to about 45°C before adding heat labile supplements.
- Aseptically dispense the desired amount of medium under a laminar airflow unit in sterile culture vessels

Storage and Shelf Life

- The plant tissue culture medium powder is extremely hygroscopic and must be stored at 2-8°C in air tight containers.
- Preferably, entire content of each package should be used immediately after opening.
- Use before the expiry date.

Central Drug House (P) Ltd. | Corp. Office : 7/28, Vardaan House, Darya Ganj, New Delhi - 110002 (INDIA), Phone : +91-11-49404040 (100 Lines) Mfg Unit : Plot No. D-2/CH/9, Dahej-2, GIDC, Dist. Bharuch - 392130 (Gujarat), E-mail : sales@cdhfinechemical.com

Plant Tissue Culture



Product Specification

cdhfinechemical.com

Precautions

- Ensure appropriate pH of the medium before addition of gelling agent as acidic pH will lead to decreased gelation resulting in semi solid flowing gel while alkaline pH will lead to formation of hardened gel.
- Use of Distilled water/Tissue culture grade water is recommended for media preparation as tap water or lower grade water may lead to salt precipitation and improper gelation.
- Avoid preparation of concentrated solutions, as it will lead to precipitation of salts.

Disclaimer

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.