

Technical Information

Carrot Callus Initiation Medium With Vitamins and 2,4-D Without Sucrose and Agar

Product Code: PT1089

Application: Carrot Callus Initiation medium has been formulated to initiate the carrot callus from its tissue. It is a nutrient blend of inorganic salts that provides all the essential macronutrients, micronutrients, vitamins and plant growth regulators required for the establishment of the species.

High amount of potassium nitrate is effective in inducing somatic embryogenesis. Sodium dihydrogen phosphate serves as a source of phosphate and promotes root development and vigorous growth. Microelements like Boron, Molybdenum, Iron, Zinc and Manganese enhance metabolism in the plants. Boron plays a key role in carbohydrate metabolism and 2,4-D aids in callus formation.

The product is plant tissue culture tested but it is the sole responsibility of the user to ensure the suitability of the medium for individual species.

Composition**

Ingredients	mg/Litre
MACROELEMENTS	
Ammonium sulphate	134.000
Calcium chloride	113.250
Magnesium sulphate	122.087
Potassium sulphate	2500.000
Sodium dihydrogen orthophosphate	130.417
MICROELEMENTS	
Boric acid	3.000
Cobalt chloride hexahydrate	0.025
Copper sulphate pentahydrate	0.025
EDTA disodium salt dehydrate	37.300
Ferrous sulphate heptahydrate	27.800
Manganese sulphate monohydrate	10.000
Molybdic acid (sodium salt)	0.213
Potassium Iodide	0.750
Zinc sulphate heptahydrate	2.000
VITAMINS	
myo-Inositol	100.000
Nicotinic acid (free acid)	1.000
Pyridoxine HCl	1.000
Thiamine hydrochloride	10.000
OTHERS	
2,4-Dichlorophenoxy acetic acid	1.000
Total	3.2 gms/litre

Material required but not provided

- Autoclaved distilled water
- Sucrose (PCT1607)
- Gelling agents like Agar (PCT1901) or CleriGel (PCT1903)
- 1N NaOH/HCl

Quality Control

Appearance

White to off-white, homogenous, free flowing powder

Solubility

3.2 gms/litre soluble in distilled water

Colour and Clarity

Colourless to light yellow, clear solution

pH at 25°C

3.50 – 4.50

Plant Tissue Culture Test

The growth promoting properties of medium is assessed by providing plant cultures with relative humidity of about 60%±2%, temperature 22°C±2°C and photoperiod of about 16:8. The plant species showed actively growing callus and shoots with no structural, necrotic and toxic deformity.

Directions

- Reconstitute medium by adding required quantity of powder in two-third of total volume with constant, gentle stirring till the medium gets completely dissolved.
- Add heat stable supplements prior to autoclaving.
- Make up the final volume with distilled water.
- Adjust the pH of the medium to 5.75 ± 0.5 using 1N NaOH/HCl.
- Add gelling agent and heat the medium to boiling till complete dissolution of gelling agent.
- Sterilize the medium by autoclaving at 15 lbs and 121°C for 15 min.
- Cool the autoclaved medium to about 45°C before adding heat labile supplements.
- Aseptically dispense the desired amount of medium under a laminar airflow unit in sterile culture vessels

Storage and Shelf Life

- The plant tissue culture medium powder is extremely hygroscopic and must be stored at 2-8°C in air tight containers.
- Preferably, entire content of each package should be used immediately after opening.
- Use before the expiry date.

Disclaimer

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

Precautions

- Ensure appropriate pH of the medium before addition of gelling agent as acidic pH will lead to decreased gelation resulting in semi solid flowing gel while alkaline pH will lead to formation of hardened gel.
- Use of Distilled water/Tissue culture grade water is recommended for media preparation as tap water or lower grade water may lead to salt precipitation and improper gelation.
- Avoid preparation of concentrated solutions, as it will lead to precipitation of salts.