

Technical Information

Litvay Medium w/ Vitamins; w/o Sucrose and Agar

Product Code: PT1042

Composition**

Ingredients	mg/Litre
Potassium nitrate	1900.00
Ammonium nitrate	1650.00
Calcium chloride.2H ₂ O	22.00
Magnesium sulphate	903.37
Potassium phosphate monobasic	340.00
Manganese sulphate.H ₂ O	21.00
Boric acid	31.00
Potassium iodide	4.15
Molybdic acid (sodium salt).2H ₂ O	1.25
Zinc sulphate.7H ₂ O	43.00
Copper sulphate.5H ₂ O	0.50
Cobalt chloride.6H ₂ O	0.125
Ferrous sulphate.7H ₂ O	27.80
EDTA disodium salt.2H ₂ O	37.26
myo - Inositol	100.00
Thiamine hydrochloride	0.10
Pyridoxine hydrochloride	0.10
Nicotinic acid (Free acid)	0.50
TOTAL	5.08 gm/litre

Principle And Interpretation

Litvay medium has been specially formulated for plant cell, tissue and organ cultures. Potassium nitrate and ammonium nitrate serves as nitrate sources. Medium does not contain sucrose and agar; hence these components have to be added to the medium before use.

Directions

Suspend 5.08 grams of dehydrated medium# in 600ml of distilled water and rinse media vial with small quantity of distilled water to remove traces of powder. Apply constant gentle stirring to the solution till the powder dissolves completely. Add desired heat stable supplements prior to autoclaving. Adjust the medium to the desired pH using 1N HCl/NaOH. Make up the final volume to 1000ml with distilled water. Sterilize the medium by autoclaving at 15 lbs or 121°C for 15 minutes. Cool the autoclaved medium to 45°C before adding the filter sterilized heat labile supplements. Dispense the desired amount of medium aseptically in sterile culture vessels.

Weight after vacuum drying to remove all water

Quality Control

Appearance	: White to off-white, homogeneous, free flowing powder.
Solubility	: 5.08 gm/litre soluble in distilled water.
Colour and Clarity	: Colourless to light yellow, clear to slight opalescent solution.
pH at 25°C	: 4.1 ±0.5 of 0.508% w/v dehydrated medium.

Cultural Response :

Cultural condition :

· Incubation period	: 5 weeks
· Relative humidity	: 60% ± 2%
· Temperature	: 22°C ± 2°C
· Photoperiod (D:N) in hours	: 16:8

Cell Line	Types Of Culture	Results
<i>Musa species</i>	Shoot culture	No structural deformity observed No necrotic tissues, Actively growing shoots, No toxicity to shoots
<i>Daucus species</i>	Callus culture	No necrotic tissues, Actively growing callus, No toxicity to callus

Storage and Shelf Life

Dehydrated plant tissue culture media powder is extremely hygroscopic and should be protected from atmospheric moisture. If possible, the entire content of each bottle should be used immediately after opening or else the unused portion should be stored in a desiccator and refrigerated at 2-8°C. Use before the expiry date.

Further Reading

1. Litvay J.D., Verma D.C. & Johnson M.A., Plant Cell Reports, (1985), 4(6), 325 - 328

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.