



Technical Information

DKW / Juglans Medium w/ Vitamins; w/o Sucrose and Agar

Product Code: PT1032

Composition**

Ingredients	mg/Litre	
Ammonium nitrate	1416.00	
Calcium chloride.2H ₂ O	149.00	
Calcium nitrate	1367.47	
Magnesium sulphate	361.38	
Potassium phosphate monobasic	265.00	
Potassium sulphate	1559.00	
Manganese sulphate.H₂O	33.50	
Boric acid	4.80	
Molybdic acid (sodium salt).2H₂O	0.39	
Zinc nitrate.6H₂O	17.00	
Copper sulphate.5H₂O	0.25	
Ferrous sulphate.7H₂O	33.80	
EDTA disodium salt.2H₂O	45.40	
myo - Inositol	100.00	
Thiamine hydrochloride	2.00	
Nicotinic acid (Free acid)	1.00	
Glycine (Free base)	2.00	
TOTAL	5.36 gm/litre	

Principle And Interpretation

DKW / Juglans medium has been specially formulated for plant cell, tissue and organ cultures. Ammonium nitrate and calcium nitrate serves as nitrate sources. Glycine serves as an amino acid source. Medium does not contain sucrose and agar; hence these components have to be added to the medium before use.

Directions

Suspend 5.36 grams of dehydrated medium# in 600ml of distilled water and rinse media vial with small quantity of distilled water to remove traces of powder. Apply constant gentle stirring to the solution till the powder dissolves completely. Add desired heat stable supplements prior to autoclaving. Adjust the medium to the desired pH using 1N HCI/NaOH.

Make up the final volume to 1000ml with distilled water. Sterilize the medium by autoclaving at 15 lbs or 121ºC for 15 minutes. Cool the autoclaved medium to 45°C before adding the filter sterilized heat labile supplements. Dispense the desired amount of medium aseptically in sterile culture vessels.

Weight after vacuum drying to remove all water





Product Specification

Quality Control

Appearance : White to off-white, homogeneous, free flowing powder.

Solubility : 5.29 gm/litre soluble in distilled water. : Colourless to light yellow, clear solution. Colour and Clarity

pH at 25°C : 4.0 ±0.5 of 0.529% w/v dehydrated macroelements powder.

<u>Cultural Response</u>:

Cultural condition:

· Incubation period : 5 weeks · Relative humidity : 60% ± 2% · Temperature : 22°C ± 2°C · Photoperiod (D:N) in hours : 16:8

Cell Line	Types Of Culture	Results	
Musa species	Shoot culture	No structural deformity observed	
		No necrotic tissues,	
		Actively growing shoots,	
		No toxicity to shoots	
Daucus species	Callus culture	No necrotic tissues,	
		Actively growing callus,	
		No toxicity to callus	

The medium is prepared as per direction. The growth promoting activity of this plant tissue culture medium is evaluated using two plant species viz. Musa species and Daucus species through three passages. Plant growth hormones (e.g. 2,4-D, NAA, Kinetin and 6-BAP) are added in suitable combinations and concentrations.]

Storage and Shelf Life

Dehydrated plant tissue culture media powder is extremely hygroscopic and should be protected from atmospheric moisture. If possible, the entire content of each bottle should be used immediately after opening or else the unused portion should be stored in a desiccator and refrigerated at 2-8°C. Use before the expiry date.

Further Reading

1. Driver J.A. & Kuniyuki A.H., Hort. Science, (1984), 19 (4), 507 - 509





Product Specification

dhfinechemical con

Disclaimer:

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performens parameters.