

Technical Information

Phyto Peptone Bromide Agar Base

Product Code: PHM1007

Application: Semi selective medium for the detection of *Xanthomonas hortorum pv. carotae* in carrot.

Composition**

| Ingredients | Grams/Litre |
|--------------------|-----------------------|
| Peptone | 10.0 |
| Boric acid | 0.3 |
| Potassium bromide | 10.0 |
| Agar | 15.0 |
| Total | 35.30 gm/liter |
| Final pH (at 25°C) | 7.4 |

**Formula adjusted standard to suit the performance parameter

Principle And Interpretation

Bacterial leaf blight of carrot (*Daucus carota subsp. sativus*) caused by *Xanthomonas hortorum pv. carotae* (also known as *Xanthomonas campestris pv. carotae*) is an important seed-borne disease found worldwide. Under conditions favourable to the bacteria this disease can cause significant yield losses. Contaminated carrot seeds are a potential primary source of inoculum and the use of *X.hortorum pv. carotae*-free seed is an important disease management strategy. Therefore, sensitive detection methods suitable for routine application are needed by many agricultural industries (1). This medium is based on the formulation of mTBM Medium as suggested by McGuire et.al 1986 (2) and serves as a semi-selective medium for the growth of *X. hortorum pv. carotae* . Peptone provides the necessary nitrogen compounds, carbon vitamins and also some trace elements necessary for the growth of plant pathogen. Boric acid serves as a selective agent. . Nystatin inhibits anti-fungal agent.

Directions

Suspend 35.3 grams in 900 ml distilled water containing 10 ml of Tween 80. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121) for 15 minutes. Dissolve 10.0 grams of skim milk powder in 100 ml of distilled water. Sterilize by autoclaving at 15 lbs pressure (121) for 5 minutes. Mix Part A and Part B, cool to 45-50°C.and aseptically add the rehydrated content of one vial of CNF supplement (PHS1006).

Mix well and pour into petri plates.

Quality Control

Appearance:

Cream to yellow coloured, homogeneous, free flowing powder.

Gelling:

Firm, comparable with 1.5% Agar gel.

Colour and Clarity of prepared medium:

Yellow coloured, opalescent gel forms in Petri plates.

Reaction:

Reaction of 3.53 % w/v aqueous solution is pH 7.4 ± 0.2 at 25°C.

Cultural Response:

Cultural characteristics observed with added CNF Supplement (PHS1006), after an incubation at 25-30°C for 5-7 days

| Organism (ATCC) | Growth | Colony Characteristics |
|---|-----------|--|
| <i>Xanthomonas hortorum pv. carotae</i> | Luxuriant | White / Yellow / White yellow, glistening round, convex with entire Margins and surrounded by a clear zone of casein |
| <i>Escherichia coli</i> (25922) | Inhibited | - |
| <i>Staphylococcus aureus</i> (25923) | Inhibited | - |
| <i>Saccharomyces cerevisiae</i> (9763) | Inhibited | - |

Storage and Shelf Life

Store below 30°C and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Further Reading

- Meijerink, G.A.A.M. and van Breukelen, E.W.M. (1995). International initiative standardizes test protocols. *Prophyta* annual 49, pp. 58-65.
- McGuire, R.G., Jones, J.B. and Sasser, M. 1986. Tween media for semiselective isolation of *Xanthomonas campestris pv vesicatoria* from soil and plant material. *Plant Dis.* 70:887-891.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.