

Technical Information

McCoy's 5A Medium

With L-Glutamine Without Folic acid ,Phenol red and Sodium bicarbonate

Product Code: AT1179F

Application:- McCoy's 5A medium was developed at Roswell Park Memorial Institute in Buffalo, New York. The first medium was developed in 1955 as the result of studies on the nutritional requirements of the Walker 256 carcinoma. The original formulation was based on the amino acids in concentrations similar to those in Eagles. medium as well as the water soluble vitamins of Medium199. Modifications to the original formulation resulted in the final version being published in 1960. The final formulation also incorporates modifications done by Iwakata and Grace and contains increased amounts of folic acid, vitamin B12, and peptone. This medium is also known to support growth of primary cultures derived from a variety of tissues.

AT1179F is McCoy's 5A medium with L-glutamine. It does not contain folic acid and phenol red. Users are advised to review the literature for recommendations regarding medium supplementation and physiological growth requirements specific for different cell lines.

Composition**

Ingredients	mg/Litre
INORGANIC SALTS	
Calcium chloride dihydrate	132.430
Magnesium sulphate anhydrous	97.720
Potassium chloride	400.000
Sodium chloride	6460.00
Sodium phosphate monobasic anhydrous	504.350
AMINO ACIDS	
Glycine	7.510
L-Arginine hydrochloride	13.360
L-Asparagine anhydrous	42.140
L-Aspartic acid	45.030
L-Cysteine dihydrochloride	19.970
L-Glutamic acid	31.500
L-Glutamine	22.070
L-Histidine hydrochloride	20.960
L-Hydroxyproline	19.670
L-Isoleucine	39.360
L-Leucine	39.360
L-Lysine hydrochloride	36.540
L-Methionine	14.920
L-Phenylalanine	16.520
L-Proline	17.270
L-Serine	26.280
L-Threonine	17.870
L-Tryptophan	3.060
L-Tyrosine disodium salt	26.100
L-Valine	17.570

VITAMINS

Ascorbic acid	0.5625
Biotin	0.200
Choline chloride	5.000
D-Ca-Pantothenate	0.200
Niacin	0.500
Niacinamide	0.500
Pyridoxal hydrochloride	0.500
Pyridoxine hydrochloride	0.500
Riboflavin	0.200
Thiamine hydrochloride	0.200
Vitamin B12	2.000
i-Inositol	36.000
p-Amino benzoic acid (PABA)	1.000

OTHERS

D-Glucose	3000.00
Glutathione reduced	0.500
Peptic digest of animal tissue	600.000

Methodology

1. Suspend 12.0gms in 900ml tissue culture grade water with constant, gentle stirring until the powder is completely dissolved. Do not heat the water.
2. Add 2.2gms of sodium bicarbonate powder (TC1230) or 29.3ml of 7.5% sodium bicarbonate solution (TCL1013) for 1 litre of medium and stir until dissolved.
3. Adjust the pH to 0.2 - 0.3 pH units below the desired pH using 1N HCl or 1N NaOH since the pH tends to rise during filtration.
4. Make up the final volume to 1000ml with tissue culture grade water.
5. Sterilize the medium immediately by filtering through a sterile membrane filter with a porosity of 0.22 micron or less, using positive pressure rather than vacuum to minimize the loss of carbon dioxide.
6. Aseptically add sterile supplements as required and dispense the desired amount of sterile medium into sterile containers.
7. Store liquid medium at 2-8°C and in dark till use.

Material required but not provided :

Tissue culture grade water (TCL1010)
 Sodium bicarbonate (TC1230)
 Sodium bicarbonate solution, 7.5% (TCL1013)
 1N Hydrochloric acid (TCL1003)
 1N Sodium hydroxide (TCL1002)

Quality Control

Foetal bovine serum (BA3112/BA12432)

Appearance

Off-white to Creamish white, homogenous powder.

Solubility

Clear solution at 12.0 gms/L.

pH without Sodium Bicarbonate

4.80 -5.40

pH with Sodium Bicarbonate

6.90 -7.50

Osmolality without Sodium Bicarbonate

240.00 -280.00

Osmolality with Sodium Bicarbonate

285.00 -325.00

Cultural Response

The growth promotion capacity of the medium is assessed qualitatively by analyzing the cells for the morphology and quantitatively by estimating the cell counts and comparing it with a control medium through minimum three subcultures.

Endotoxin content

NMT 1EU/ml

Storage and Shelf Life

1. All the powdered media and prepared liquid culture media should be stored at 2-8°C. Use before the expiry date. In spite of above recommended storage condition, certain powdered medium may show some signs of deterioration /degradation in certain instances. This can be indicated by change in colour, change in appearance and presence of particulate matter and haziness after dissolution.
2. Preparation of concentrated medium is not recommended since free base amino acids and salt complexes having low solubility may precipitate in concentrated medium.
3. pH and sodium bicarbonate concentration of the prepared medium are critical factors affecting cell growth. This is also influenced by amount of medium and volume of culture vessel used (surface to volume ratio). For example, in large bottles, such as Roux bottles pH tends to rise perceptibly as significant volume of carbon dioxide is released. Therefore, optimal conditions of pH, sodium bicarbonate concentration, surface to volume ratio must be determined for each cell type. We recommend stringent monitoring of pH. If needed, pH can be adjusted by using sterile 1N HCl or 1N NaOH or by bubbling in carbon dioxide.
4. If required, supplements can be added to the medium prior to or after filter sterilization observing sterility precautions. Shelf life of the medium will depend on the nature of supplement added to the medium.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.