



Product Specification

Technical Information

Dulbecco's Modified Eagle Medium (DMEM) With 4.5gms Glucose per litre and Sodium pyruvate Without L-Glutamine and Sodium bicarbonate

Product Code: AT1007A

Application: - Dulbecco's Modified Eagle Medium (DMEM) is one of the most widely used modification of Eagle's medium. DMEM is a modification of Basal Medium Eagle (BME) that contains four fold concentration of amino acids and vitamins. Additionally, the formulation also includes glycine, serine and ferric nitrate. The original formulation contains 1000mgs/L of glucose and was originally used to culture embryonic mouse cells. DMEM high glucose is a further modification of original DMEM and contains 4500mgs glucose per litre. The additional glucose has proved to be useful in cultivating various other cell lines including primary cultures of mouse and chicken cells as well as various normal and transformed cell lines. AT1007A is Dulbecco's Modified Eagle Medium with 4.5gm glucose per litre and sodium pyruvate. It does not contain L-Glutamine. Users are advised to review the literature for recommendations regarding medium supplementation and physiological growth requirements specific for different cell lines.

Composition**

Ingredients	Gms / Litre
INORGANIC SALTS	
Calcium chloride dihydrate	265.000
Ferric nitrate nonahydrate	0.100
Magnesium sulphate anhydrous	97.720
Potassium chloride	400.000
Sodium chloride	6400.000
Sodium dihydrogen phosphate anhydrous	109.000
AMINO ACIDS	
Glycine	30.000
L-Arginine hydrochloride	84.000
L-Cystine dihydrochloride	62.570
L-Histidine hydrochloride monohydrate	42.000
L-Isoleucine	105.000
L-Leucine	105.000
L-Lysine hydrochloride	146.000
L-Methionine	30.000
L-Phenylalanine	66.000
L-Serine	42.000
L-Threonine	95.000
L-Tryptophan	16.000
L-Tyrosine disodium salt	103.790
L-Valine 	94.000
VITAMINS	
Choline chloride	4.000
D-Ca-Pantothenate	4.000
Folic acid	4.000
Nicotinamide	4.000





Product Specification

dhfinechemical.com

Pyridoxal hydrochloride	4.000
Riboflavin	0.400
Thiamine hydrochloride	4 .000
i-Inositol	7.200

OTHERS

D-Glucose 4500.000
Phenol red sodium salt 16.000
Sodium pyruvate 110.000

Methodology

- Suspend 13.0gms in 900ml tissue culture grade water with constant, gentle stirring until the powder is completely dissolved. Do not heat the water
 Add 3.7gms of sodium bicarbonate powder (TC1230) or 49.3ml of 7.5% sodium bicarbonate solution (TCL1013) and 0.584gms of L-Glutamine powder
 (TC1243) or 20ml of 200mM L-Glutamine solution (TCL1012)for 1 litre of medium and stir until dissolved.
- 3. Adjust the pH to 0.2-0.3 pH units below the desired pH using 1N HCl o since the pH tends to rise during filtration
- 4. Make up the final volume to 1000ml with tissue culture grade water
- 5. Sterilize the medium immediately by filtering through a sterile membrane filter with a porosity of 0.22 micron or less, using positive pressure rather than vacuum to minimize the loss of carbon dioxide
- 6. Aseptically add sterile supplements as required and dispense the desired amount of sterile medium into sterile containers
- 7. Store liquid medium at 2-8°C and in dark till use

Material required but not provided

Tissue culture grade water (TCL1010)

Sodium bicarbonate powder (TC1230)

Sodium bicarbonate solution, 7.5% (TCL1013)

1N Hydrochloric acid (TCL1003)

1N Sodium hydroxide (TCL1002)

L-Glutamine powder (TC1243)

L-Glutamine solution 200mM (TCL1012)

Fetal bovine serum (BA3112/BA30432)

Quality Control

Appearance

Off-white to Creamish white, homogenous powder.

Solubility

Clear solution at 13.0gms/L

pH without Sodium Bicarbonate

5.90-6.50

pH with Sodium Bicarbonate

7.40 -8.00

Osmolality without Sodium Bicarbonate

250.00 -290.00

Osmolality with Sodium Bicarbonate

330.00 -370.00

Cultural Response

The growth promotion capacity of the medium is assessed qualitatively by analyzing the cells for the morphology and quantitatively by estimating the cell counts and comparing it with a control medium through minimum three subcultures.

Endotoxin Content

NMT 5EU/ml





Product Specification

Storage and Shelf Life

1. All the powdered media and prepared liquid culture media should be stored at 2-8°C. Use before the expiry date. In spite of above recommended storage condition, certain powdered medium may show some signs of deterioration /degradation in certain instances. This can be indicated by change in colour, change in appearance and presence of particulate matter and haziness after dissolution.

2. Preparation of concentrated medium is not recommended since free base amino acids and salt complexes having low solubility may precipitate in concentrated medium.

3. pH and sodium bicarbonate concentration of the prepared medium are critical factors affecting cell growth. This is also influenced by amount of medium and volume of culture vessel used (surface to volume ratio). For example, in large bottles, such as Roux bottles pH tends to rise perceptibly as significant volume of carbon dioxide is released. Therefore, optimal conditions of pH sodium bicarbonate concentration, surface to volume ratio must be determined for each cell type. We recommend stringent monitoring of pH. If needed, pH can be adjusted by using sterile 1N HCl or 1N NaOH or by bubbling in carbon dioxide.

4. If required, supplements can be added to the medium prior to or after filter sterilization observing sterility precautions. Shelf life of the medium will depend on the nature of supplement added to the medium.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate.
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performens parameters.