

## **Technical Information**

# Dulbecco's Modified Eagle Medium/ Nutrient Mixture F12 Ham (DMEM/F12, 1:1 mixture) With L-Glutamine

Without HEPES buffer, Trace elements and Sodium bicarbonate

Product Code: AT1155

Application:- Dulbecco's Modified Eagle Medium / Nutrient Mixture F12 Ham DMEM/F12, 1:1 mixture) was originally formulated for rat neuroblastoma cells and MDCK cells. The mixture is extremely nutritious and supports growth of a wide variety of cells including certain epithelial, endothelial and granulosa cells.

AT1155 is DMEM/ Nutrient Mixture F-12 Ham with L- glutamine. It does not contain HEPES buffer, phenol red and trace elements..Users are advised to review the literature for recommendations regarding medium supplementation and physiological growth requirements specific for different cell lines.

Composition**		
Ingredients	mg/Litre	
INORGANIC SALTS		
Calcium chloride dihydrate	154.500	
Copper sulphate pentahydrate	0.0013	
Disodium hydrogen phosphate, anhydrous	71.020	
Ferric nitrate nonahydrate	0.050	
Ferrous sulphate heptahydrate	0.417	
Magnesium chloride anhydrous	61.200	
Magnesium sulphate anhydrous	48.840	
Potassium chloride	311.800	
Sodium chloride	6996.00	
Sodium dihydrogen phosphate anhydrous	54.300	
Zinc sulphate heptahydrate	0.432	
AMINO ACIDS		
Glycine	18.750	
L-Alanine	4.450	
L-Arginine hydrochloride	147.500	
L-Asparagine monohydrate	7.500	
L-Aspartic acid	6.650	
L-Cysteine dihydrochloride	17.560	
L-Cystine hydrochloride monohydrate	31.290	
L-Glutamic acid	7.350	
L-Glutamine	365.00	
L-Histidine hydrochloride monohydrate	31.480	
L-Isoleucine	54.470	
L-Leucine	59.050	
L-Lysine hydrochloride	91.250	
L-Methionine	17.240	
L Wethornie	17.270	





## **Product Specification**

cdhfinechemical.com

L-Phenylalanine	35.480
Prolin	17.250
L-Serine	26.250
L-Threonine	53.450
L-Tryptophan	9.020
L-Tyrosine disodium salt	48.100
L-Valine	52.850
VITAMINS	
Choline chloride	8.980
D-Biotin	0.0035
D-Pantothenic acid (hemicalcium)	2.240
Folic acid	2.660
Niacinamide	2.020
Pyridoxal hydrochloride	2.000
Pyridoxine hydrochloride	0.031
Riboflavin	0.219
Thiamine hydrochloride	2.170
Vitamin B12	0.680
myo-Inositol	12.600
OTHERS	
D-Glucose	3151.000
DL-Thioctic Acid	0.105
Hypoxanthine	2.400
Linoleic acid	0.042
Phenol red sodium salt	8.630
Putrescine hydrochloride	0.081
Sodium pyruvate	110.000
Thymidine	0.365

## Methodology

- 1. Suspend 12.1gms in 900ml tissue culture grade water with constant, gentle stirring until the powder is completely dissolved. Do not heat the water.
- 2. Add 1.2gms of sodium bicarbonate powder (TC1230) or 16ml of 7.5% Sodium bicarbonate solution (TCL1013) for1 litre of medium and stir until dissolved.
- 3. Adjust the pH to 0.2 0.3 pH units below the desired pH using 1N HCl or 1N NaOH since the pH tends to rise during filtration.
- 4. Make up the final volume to 1000ml with tissue culture grade water.
- 5. Sterilize the medium immediately by filtering through a sterile membrane filter with a porosity of 0.22 micron or less, using positive pressure rather than vacuum to minimize the loss of carbon dioxide.
- 6. Aseptically add sterile supplements as required and dispense the desired amount of sterile medium into sterile containers.
- 7. Store liquid medium at 2-8°C and in dark till use.

## Material required but not provided:

Tissue culture grade water (TCL1010)

Sodium bicarbonate powder (TC1230)

Sodium bicarbonate solution, 7.5% (TCL1013)

1N Hydrochloric acid (TCL1003)

1N Sodium hydroxide (TCL1002)

Foetal bovine serum (BA3112/BA12432)

## **Quality Control**

#### Appearance

Off-white to Creamish white, homogenous powder.

#### Solubility

Clear solution at 12.1 gms/L.

#### pH without Sodium Bicarbonate

6.00 -6.60

#### pH with Sodium Bicarbonate

7.20 -7.80

#### Osmolality without Sodium Bicarbonate

260.00 -300.00

#### Osmolality with Sodium Bicarbonate

285.00 -325.00

#### **Cultural Response**

The growth promotion capacity of the medium is assessed qualitatively by analyzing the cells for the morphology and quantitatively by estimating the cell counts and comparing itwith a control medium through minimum three subcultures.

#### **Endotoxin Content**

NMT 5EU/ml

## Storage and Shelf Life

- 1. All the powdered media and prepared liquid culture media should be stored at 2-8°C. Use before the expiry date. In spite of above recommended storage condition, certain powdered medium may show some signs of deterioration /degradation in certain instances. This can be indicated by change in colour, change in appearance and presence of particulate matter and haziness after dissolution.
- 2. Preparation of concentrated medium is not recommended since free base amino acids and salt complexes having low solubility may precipitate in concentrated medium.
- 3. pH and sodium bicarbonate concentration of the prepared medium are critical factors affecting cell growth. This is also influenced by amount of medium and volume of culturevessel used (surface to volume ratio). For example, in large bottles, such as Roux bottles pH tends to rise perceptibly as significant volume of carbon dioxide is released. Therefore, optimal conditions of pH, sodium bicarbonate concentration, surface to volume ratio must be determined for each cell type. We recommend stringent monitoring of pH. If needed, pH can be adjusted by using sterile 1N HCl or 1N NaOH or by bubbling in carbon dioxide.
- 4. If required, supplements can be added to the medium prior to or after filter sterilization observing sterility precautions. Shelf life of the medium will depend on the nature of supplement added to the medium.

#### Disclaimer:

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.





# **Product Specification**

cdhfinechemical.com