



# **Product Specification**

cdhfinechemical.com

# **Technical Information**

## **Minimum Essential Medium Eagle (MEM)**

With Earle's salts, L-Glutamine, NEAA and Sodium bicarbonate Hybridoma Tested 1X Liquid Cell Culture Medium

### Product Code: AL1047H

**Application:** Minimum Essential Medium (MEM) is a modification of Basal Medium Eagle (BME). It was developed by Harry Eagle to meet the specific nutritional requirements of certain subtypes of HeLa cells and normal mammalian fibroblasts. MEM includes higher concentration of amino acids so as to closely approximate the protein composition of cultured mammalian cells. MEM can be used either with Earle's salts or Hank's salts and can also be additionally supplemented with non-essential amino acids (NEAA). This medium can be further modified by eliminating calcium to facilitate growth of cells in suspension cultures.

AL1047H is Minimum Essential Medium Eagle with Earle's salts, L-glutamine, sodium bicarbonate and nonessential amino acids. This medium is also hybridoma tested. Users are advised to review the literature for recommendations regarding medium supplementation and physiological growth requirements specific for different cell lines.

| <b>O</b> |     | _ • • • |      | **           |
|----------|-----|---------|------|--------------|
| Con      | ากก | (dilli) | Λh   | <b>TO TO</b> |
|          |     | P117    | VIII |              |

| Ingredients                           | mg/Litre |
|---------------------------------------|----------|
| INORGANIC SALTS                       |          |
| Calcium chloride dihydrate            | 265.000  |
| Magnesium sulphate anhydrous          | 97.720   |
| Potassium chloride                    | 400.000  |
| Sodium bicarbonate                    | 2200.000 |
| Sodium chloride                       | 6800.000 |
| Sodium dihydrogen phosphate anhydrous | 122.000  |
| AMINO ACIDS                           |          |
| Glycine                               | 7.500    |
| L-Alanine                             | 8.900    |
| L-Arginine hydrochloride              | 126.000  |
| L-Asparagine monohydrate              | 15.000   |
| L-Aspartic acid                       | 13.300   |
| L-Cystinedihydrochloride              | 31.300   |
| L-Glutamic acid                       | 14.700   |
| L-Glutamine                           | 292.000  |
| L-Histidine hydrochloride monohydrate | 42.000   |
| L-Isoleucine                          | 52.000   |
| L-Leucine                             | 52.000   |
| L-Lysine hydrochloride                | 72.500   |
| L-Methionine                          | 15.000   |
| L-Phenylalanine                       | 32.000   |
| L-Proline                             | 11.500   |
| L-Serine                              | 10.500   |
| L-Threonine                           | 48.000   |
|                                       | 10.000   |
| L-Tryptophan                          | 10.000   |





# **Product Specification**

cdhfinechemical.com

| -Valine                | 46.000   |  |
|------------------------|----------|--|
| /ITAMINS               |          |  |
| Choline chloride       | 1.000    |  |
| )-Ca-Pantothenate      | 1.000    |  |
| olic acid              | 1.000    |  |
| licotinamide           | 1.000    |  |
| yridoxal hydrochloride | 1.000    |  |
| Riboflavin             | 0.100    |  |
| hiamine hydrochloride  | 1.000    |  |
| Inositol               | 2.000    |  |
| OTHERS                 |          |  |
| )-Glucose              | 1000.000 |  |
| henol red sodium salt  | 11.000   |  |

## **Quality Control**

#### Appearance

Orangish red colored, clear solution.

#### рΗ

7.00 -7.60

#### Osmolality in mOsm/Kg H<sub>2</sub>O

295.00 -335.00

#### Sterility

No bacterial or fungal growth is observed after 14 days of incubation, as per USP specification.

#### Cultural Response

The growth promotion capacity of the medium is assessed qualitatively by analyzing the cells for the morphology and quantitatively by estimating the cell counts and comparing it with a control medium through minimum three subcultures.

#### **Endotoxin Content**

NMT 5EU/ml

## Storage and Shelf Life

Store at 2-8°C away from bright light. Shelf life is 12 months. Use before expiry date given on the product label.

### Disclaimer:

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.