

## Technical Information

### SOC Growth Medium

#### Product Code: G1015

SOC Growth Medium is a special medium for incubating competent cells immediately after transformation to allow expression of transferred resistance genes before exposing cells to selective conditions.

#### Composition\*\*

Ingredients	Grams/Litre
Tryptone	20.00
Yeast extract	5.00
Sodium chloride	0.50
MgSO <sub>4</sub> ·7H <sub>2</sub> O	5.00
Glucose	3.6

\*\* Formula adjusted, standardized to suit performance parameters

#### Methodology

Suspend 31.54 grams of dehydrated media in 1000 ml distilled water. Heat if necessary to dissolve the medium completely. Dispense as desired and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

#### Principle and Interpretation

SOC Growth Medium is a special medium for incubating competent cells immediately after transformation to allow expression of transferred resistance genes before exposing cells to selective conditions (1). *E. coli* cells are first made competent during transformation where perforations are made in the bacterial cells so that the foreign DNA can penetrate the cells. To repair cells from this damage SOC medium is used. This medium is same as SOB medium except glucose is added as the carbon source which repairs the perforations of the *E. coli* cells. Furthermore, the transferred resistance genes are expressed in this medium before exposure to selective conditions (1). Tryptone provides nitrogen, amino acids and other growth factors which permit the cells to go through the stress of transformation. Vitamins and trace elements are contained in Yeast Extract. Sodium chloride provides essential ions for transport and osmotic balance. Magnesium sulfate provides magnesium ions which are required in a variety of enzymatic reactions, including DNA replication

#### Quality control

##### Appearance of Powder :

Light yellow coloured, homogeneous, free flowing powder.

##### Colour and Clarity :

Light yellow coloured, clear solution without any precipitate.

##### Cultural Response :

Cultural characteristics observed after an incubation at 35-37°C for 18 - 48 hours.

##### Organisms (ATCC)

*Escherichia coli* ATCC 23724

*Escherichia coli* ATCC 25922

*Escherichia coli* MTCC 1652

##### Growth

good-luxuriant

good-luxuriant

good-luxuriant

#### Storage and Shelf Life

Store below 30°C and the prepared medium at 2 - 8°C. Use before expiry date on the label.

#### Reference:

(1) Sambrook J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2<sup>nd</sup> ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.

## **Disclaimer :**

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.