Molecular Biology Growth Media

Technical Information

SD Growth Medium w/o TRP

Product Code: G1066

SD Growth Medium w/o TRP is a synthetic defined media for the growth of Saccharomyces cerevisiae.

Composition**

Composition**		
Ingredients	Grams/Litre	
Potassium dihyrogen phosphate	1.00	
Magnesium sulphate	0.50	
Sodium chloride	0.10	
Calcium chloride	0.10	
Biotin	0.002 gm	
Calcium pantothenate	0.4 mg	
Folic acid	0.002 mg	
Inositol	2.00 mg	
Niacin	0.4 mg	
PABA	0.2 mg	
Pyridoxin, HCl	0.4 mg	
Riboflavin	0.2 mg	
Thiamine HCl	0.4 mg	
Boric acid	0.5 mg	
Copper sulphate	0.04 mg	
Potassium iodide	0.1 mg	
Ferric chloride	0.2 mg	
Manganese sulphate	0.4 mg	
Sodium molybdate	0.2 mg	
Zinc sulphate	0.4 mg	
Ammonium sulphate	5.00	
Dextrose	20.00	
Adenine	0.010	
L-Arginine HCl	0.050	
L-Aspartic acid	0.080	
L-Histidine HCl	0.020	
L-Isoleucine	0.050	
L-Leucine	0.100	
L-Lysine HCl	0.050	
L-Methionine	0.020	
L-Phenylalanine	0.050	
L-Threonine	0.100	
L-Tryptophan	0.050	
L-Tyrosine	0.050	
Uracil	0.020	
L-Valine	0.140	
** Formula adjusted, standardized to suit	performance parameters	

Methodology

Suspend 27.44 grams in 1000 ml distilled water. Sterilize by autoclaving at 10 lbs pressure (115°C) for 20 minutes. Mix well and dispense as desired.



Molecular Biology Growth Media

Principle and Interpretation

SD Growth Medium w/o TRP is a synthetic defined agar media for the selective growth of *Saccharomyces cerevisiae*. Synthetically Defined media known as Yeast Nitrogen Base Media for the growth of Yeast cells were first cited by Wickerham (1, 2). SD Growth Medium w/o TRP includes a yeast nitrogen base along with ammonium sulfate, and dextrose as the carbon source, which is further supplemented with various amino acids except tryptophan which makes it a dropout growth medium for yeast cells. A tryptophan auxotrophic yeast mutant cannot grow on this media but a wild-type or a tryptophan prototrophic yeast strain can grow. The tryptophan auxotroph has a mutation in a gene (e.g. *TRP1*) of the tryptophan synthesis pathway and this mutant strain will grow in this medium if tryptophan is supplied from outside e.g. from a plasmid which contains *TRP1* gene (3). For this purpose, a *trp1* mutant strain of *S. cerevisiae* is transformed with a *TRP1* containing plasmid and the transformants can be selected by growing the cells on SD Growth Media w/o TRP. Hence this medium is very useful in molecular genetics.

Quality control

Appearance of Powder:

White to cream colored, homogeneous, free flowing powder.

Colour and Clarity:

Light yellow coloured, clear solution without any precipitate.

Cultural Response:

Cultural characteristics observed after an incubation at 25-30°C for 18 - 48 hours.

Organisms (ATCC) Growth
Saccharomyces cerevisiae ATCC 9763 good-luxuriant

Storage and Shelf Life

Upon receipt, store at 2 - 8°C. Use before expiry date on the label.

Reference

- 1. Wickerham L. J., 1951, U.S. Dept. Agric. Tech. Bull. No. 1029
- 2. Wickerham L. J., 1946, J. Bacteriol., 52:293
- 3. Kaiser, C., et al. Methods in Yeast Genetics Cold Spring Harbor, (1994)

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is
 accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.