

Technical Information

SDA Growth Medium w/o HIS-URA

Product Code: G1076

SD Growth Medium w/o HIS-URA is a synthetic defined media for the growth of *Saccharomyces cerevisiae*.

Composition**

Ingredients	Grams/Litre
Potassium dihydrogen phosphate	1.00
Magnesium sulphate	0.50
Sodium chloride	0.10
Calcium chloride	0.10
Biotin	0.002 gm
Calcium pantothenate	0.4 mg
Folic acid	0.002 mg
Inositol	2.00 mg
Niacin	0.4 mg
PABA	0.2 mg
Pyridoxin, HCl	0.4 mg
Riboflavin	0.2 mg
Thiamine HCl	0.4 mg
Boric acid	0.5 mg
Copper sulphate	0.04 mg
Potassium iodide	0.1 mg
Ferric chloride	0.2 mg
Manganese sulphate	0.4 mg
Sodium molybdate	0.2 mg
Zinc sulphate	0.4 mg
Ammonium sulphate	5.00
Dextrose	20.00
Adenine	0.010
L-Arginine HCl	0.050
L-Aspartic acid	0.080
L-Isoleucine	0.050
L-Leucine	0.100
L-Lysine HCl	0.050
L-Methionine	0.020
L-Phenylalanine	0.050
L-Threonine	0.100
L-Tryptophan	0.050
L-Tyrosine	0.050
L-Valine	0.140
Agar	15.00

** Formula adjusted, standardized to suit performance parameters

Methodology

Suspend 42.45 grams in 1000 ml distilled water. Sterilize by autoclaving at 10 lbs pressure (115°C) for 20 minutes. Mix well and dispense as desired.

Principle and Interpretation

SDA Growth Medium w/o HIS-URA is a synthetic defined agar media for the selective growth of *Saccharomyces cerevisiae*. Synthetically Defined media known as Yeast Nitrogen Base Media for the growth of Yeast cells were first cited by Wickerham. SDA Growth Medium w/o HIS-URA includes a yeast nitrogen base along with ammonium sulfate, and dextrose as the carbon source, which is further supplemented with various amino acids except histidine and uracil which makes it a dropout growth medium for yeast cells (1, 2). A histidine and uracil auxotrophic yeast mutant cannot grow on this media but a wild-type or a histidine and uracil prototrophic yeast strain can grow. The histidine and uracil auxotroph has mutation in the genes (e.g. *HIS3* and *URA3*) of the histidine as well as uracil synthesis pathway and this mutant strain will grow in this medium if histidine and uracil are supplied from outside e.g. from plasmids which contain *HIS3* and *URA3* genes (3). For this purpose, a *his3ura3* mutant strain of *S. cerevisiae* is transformed with *HIS3* and *URA3* containing plasmids and the transformants can be selected by growing the cells on SDA Growth Media w/o HIS-URA. Hence this medium is very useful in molecular genetics.

Quality control

Appearance of Powder :

Cream to white coloured, homogeneous, free flowing powder.

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity :

Light yellow coloured, clear to slightly opalescent gel forms in Petri plates.

Cultural Response :

Cultural characteristics observed after incubation at 25-30°C for 18 - 48 hours.

Organisms (ATCC)

Saccharomyces cerevisiae ATCC 9763

Growth

good-luxuriant

Storage and Shelf Life

- Upon receipt, store at 2 - 8°C. Use before expiry date on the label.

Reference

1. Adams, A., D. E. Gottschling, C. A. Kaiser, and T. Stearns. 1997. Methods in yeast genetics: A Cold Spring Harbor Laboratory Course Manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
2. Burke, D., Dawson, D., and T. Stearns. 2000. Method in yeast genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.