

## Technical Information

### MUG MacConkey Agar

**Product Code: DM 2080**

**Application:** - MUG MacConkey Agar is recommended for the selective isolation and detection of lactose fermenting coliform organisms by a fluorogenic procedure.

### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	20.000
Lactose	10.000
Bile salts mixture	1.500
Sodium chloride	5.000
Neutral red	0.030
Crystal violet	0.001
4-Methylumbelliferyl $\beta$ -D-glucuronide (MUG)	0.100
Agar	15.000
Final pH ( at 25°C)	7.1 $\pm$ 0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Principle & Interpretation

MUG MacConkey Agar is employed for the cultivation of enteric bacteria and in differentiation of lactose fermenters and non- fermenters. The medium contains bile salts to prevent non-intestinal bacteria and lactose with neutral red indicator to distinguish the lactose-fermenting coliforms from the lactose-non-fermenting *Salmonella* and dysentery groups (1). MUG MacConkey Agar is based on the modification of MacConkey medium as per Trepeta and Edberg (2). It is used for the selective isolation and detection of lactose fermenting coliform organisms by a fluorogenic procedure. MUG MacConkey Agar helps to detect the presence of an enzyme  $\beta$ -glucuronidase and thereby rapidly identifying *Escherichia coli* in mixed clinical specimens (3).

Peptic digest of animal tissue provides essential nitrogen compounds for the growth of coliforms. Lactose acts as a the fermentable carbohydrates source. Bile salts and crystal violet enhance the growth of gram-positive bacteria.

Neutral red acts as pH indicator. MUG is cleaved by the enzyme  $\beta$ -glucuronidase to release an end product 4- methylumbelliferone which produces a visible greenish-blue fluorescence under long wave ultra-violet light (366nm). The plates are exposed to ammonia fumes to increase fluorescence as suggested by Freir and Hartman (5).

The medium can be directly inoculated with the test specimen by streaking.

### Methodology

Suspend 51.63 grams of dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat to boil to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Shake well before pouring.

### Quality Control

#### Appearance

Light yellow to pink homogeneous free flowing powder

#### Gelling

Firm, comparable with 1.5% Agar gel



Dehydrated Culture Media  
Bases / Media Supplements

#### Colour and Clarity

Red with purple tinge clear to slightly opalescent gel forms in Petri plates

#### Reaction

Reaction of 5.16% w/v aqueous solution at 25°C. pH : 7.1±0.2

#### pH Range

6.90-7.30

#### Cultural Response

DM 2080: Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 24 hours.

Organism	Inoculum (CFU)	Growth	Recovery	*Fluorescence under uv
<b>Cultural Response</b> <i>Enterobacter aerogenes</i> ATCC 13048	50-100	luxuriant	>=50%	negative
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	>=50%	positive

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and prepared medium below 2-8°C. Use before expiry period on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Cruickshank R., Duguid J. P., Marmion B. P., Swain R. H. A., 1975, Medical Microbiology, Churchill Livingstone
2. Trepeta R. W. and Edberg S. C., 1984, J. Clin. Microbiol., 19 (2) :172.
3. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
4. Maddocks J. L. and Greenan M. J. (1975) J. Clin. Pathol. 28. 686-687.
5. Freir T. A. and Hartman P. A. (1987) Appl. Env. Microbiol. 53. 1246-1250.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.

