

Technical Information

Toluidine Blue DNA Agar, Modified

Product Code: DM 1613F

Application: - Toluidine Blue DNA Agar, Modified is used for detection of thermostable deoxyribonuclease activity and establish speciation of *S. aureus* in contaminated foods in accordance with FDA BAM, 1998.

Composition**

Ingredients	Gms / Litre
Deoxyriboneuclic acid (DNA)	0.300
Calcium chloride (anhydrous)	0.0011
Sodium chloride	10.000
Toluidine blue	0.083
Tris (hydroxy methyl) amino methane	6.100
Agar	10.000
Final pH (at 25°C)	9.0±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Toluidine Blue DNA Agar, Modified is used for detection of thermostable deoxyribonuclease activity and establish speciation of *S. aureus* in contaminated foods in accordance with FDA BAM, 1998. *Staphylococcus* is a Gram-positive bacteria which includes several species that can cause a wide variety of infections in humans and other animals through infection or the production of toxins. Staphylococcal intoxication worldwide stands out as one of the main food-borne diseases, with a frequency in incidents of microbiological origin second only to salmonellosis(1). Foods that lack hygienic conditions and that are improperly stored can aid in the Staphylococcal growth and toxin production. Since the infection is caused due to toxin, Staphylococcal food poisoning is also called as 'food intoxication' (2). Toluidine Blue DNA Agar, Modified (DM 1613F) is recommended for detection of thermostable deoxyribonuclease activity and establish speciation of *S. aureus* in contaminated foods in accordance with FDA BAM, 1998(3) with a slight modification in concentration of calcium chloride and toluidine blue. DNA in the medium enables the detection of DNase activity by getting depolymerized and forming a clear zone around the microbial growth. Inclusion of toluidine blue aids in detection of DNase activity by the production of a visible bright rose- pink coloured reaction due to its metachromatic properties.

Total plate count of the suspected sample is carried out using Baird Parker Agar (DM 1043). Speciation of the organism can be confirmed using Toluidine Blue DNA Agar, Modified. Like the coagulase test, this is a highly specific test for the confirmation of the organism. The test is carried out in petriplates or microslides containing the media, prepared by spreading appropriate amount (3ml if slide) of the Toluidine Blue DNA Agar, Modified medium. 10 to 12 wells of 1mm diameter are prepared in the plates/slides after solidification. Add 0.01 ml of the broth culture into these wells after heating in a boiling water bath for 15min. Incubate (in a moist chamber) for 4 hrs at 35°C. Development of bright pink halo extending at least 1 mm from periphery of well indicates a positive reaction. Tris amino methane forms the buffering system. Sodium chloride and calcium chloride provide the ions and also maintains osmotic equilibrium.

Methodology

Suspend 26.48 grams of dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat to boil to dissolve the medium completely and continue to boil for 1 to 2 minutes. Sterilization is not necessary. Dispense into sterile Petri plates.

Quality Control

Appearance

Light yellow to light grey homogeneous free flowing powder

Gelling

Firm, comparable with 1.0% Agar gel.

Colour and Clarity

Blue coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 2.65% w/v aqueous solution at 25°C. pH : 9.0±0.2

pH Range

8.80-9.20

Cultural Response

DM 1613F: 18 hrs old BHI broth culture is heated in boiling water bath for 15 minutes and studied for thermonuclease activity. 5 mm cut wells are cut in agar plates and is filled with 25-30 µl of this culture and incubated at 35-37°C for 4 hrs (or it can also be incubated at 50°C for 2 hrs) and observed for results.

Organism

Staphylococcus aureus ATCC 12600

Staphylococcus epidermidis ATCC 14990

DNase activity

positive reaction, pink haloes extending 1mm beyond the well

negative reaction

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Bautista, L., Gaya, P., Medina, M. and Nuñez, M. 1988. Applied and Environmental Microbiology, 54(2): 566-569.
2. Murray, P. R., Baron, E. J., Jorgensen, J. H., Tenover, M. C. and Tenover, R. H. 2003. Manual of Clinical Microbiology. 8 ed. Washington, D.C: ASM.
3. FDA, U.S. 1998. Bacteriological Analytical Manual. 8 ed. Gaithersburg, MD: AOAC International.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specification for identity and performance parameters.