Dehydrated Culture Media
Bases / Media Supplements

# Technical Information

## Luria Bertani Broth, Miller pH 7.0

## Product Code: DM 3119

**Application: -** Recommended for the cultivation and maintenance of recombinant strains of Escherichia coli and may be used for routine cultivation of not particularly fastidious microorganisms.

## Composition**

| Ingredients | Gms / Litre |
|---|---|
| MG Tryptone P# | 10.000 |
| Yeast Autolysate | 5.000 |
| Sodium chlorid | 10.000 |
| Final pH (at 25°C) | 7.0 ± 0.2 |

**Formula adjusted, standardized to suit performance parameters

## Principle & Interpretation

Luria Bertani Broth, Miller is slightly different with double amount of sodium chloride as compared to original media described by Lennox for cultivation and maintenance of recombinant strains of Escherichia coli (1,2,3). The media is nutritionally rich for the growth of pure cultures of recombinant strains. Strains derived from Escherichia coli K12 are deficient in Vitamin B synthesis which are further modified by specific mutation to create auxotrophic strains and are therefore unable to grow on nutritionally deficient media. MG Tryptone P# provides peptides while Vitamin B complex is provided by yeast Autolysate. Sodium chloride provides sodium ions for membrane transport and also maintains the osmotic equilibrium of the medium.

**Type of specimen**

Isolated Microorganisms

**Specimen Collection and Handling**

For Isolated Microorganisms samples follow appropriate techniques for handling specimens as per established guidelines (4,5). After use, contaminated materials must be sterilized by autoclaving before discarding.

**Warning and Precautions**

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

**Limitations**

1. Further biochemical testing is required for complete identification.

**Performance and Evaluation**

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

## Methodology

Suspend 25.0 grams in 1000 ml purified/distilled water. Heat if necessary to dissolve the medium completely. Mix well. Dispense in tubes or flasks as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

## Quality Control

**Appearance**

Cream to yellow homogeneous free flowing powder

**Colour and Clarity of prepared medium**

Yellow to amber coloured, clear solution in tubes.

**Reaction**

Reaction of 2.5% w/v aqueous solution at 25°C. pH : 7.0 ± 0.2

**pH**

6.80-7.20

**Cultural Response**

Cultural characteristics observed after an incubation at 35-37°C for 16-24 hours.

| Organism | Inoculum (CFU) | Growth |
|---|---|---|
| Escherichia coli ATCC 23724 | 50 -100 | luxuriant |
| Escherichia coli ATCC 25922 (00013*) | 50 -100 | luxuriant |
| Escherichia coli DH5 alpha MTCC 165 | 50 -100 | luxuriant |

Key : *Corresponding WDCM numbers.

## Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 15-25°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

## Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (4,5).

## Further Reading

1. Lennox E.S./ 1955, Transduction of Linked Genetic Characters of the host by bacteriophage P1., Virology, 1:190.
2. Miller, J.H.: Experiments in molecular genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York (1972).
3. Sambrook, J., et al., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York (1989).
4. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
5. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specificatons for identity and performens parameters.