

## Technical Information

### Crystal Violet Tetrazolium Agar Base

**Product Code: DM 1586**

**Application:** - Crystal Violet Tetrazolium Agar is used for detection of gram-negative psychrotrophic bacteria causing food spoilage.

#### Composition\*\*

Ingredients	Gms / Litre
Casein enzymic hydrolysate	5.000
Yeast extract	2.500
Dextrose	1.000
Crystal violet	0.001
Agar	15.000
Final pH ( at 25°C)	7.0±0.2

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

Microorganisms which are able to grow at refrigeration temperatures are usually referred to as psychrophilic. Species of *Achromobacter*, *Alcaligenes*, *Flavobacterium* and *Pseudomonas* are included among the psychrotrophic bacteria as these organisms are able to grow relatively rapidly at commercial refrigeration temperatures (1). Many psychrotrophic microorganisms when present in large numbers can cause a variety of off-flavors as well as physical defects in foods. Their growth rate is highly dependent on temperature, and therefore, if the temperature is reduced, their growth rate is also slowed down. Thus the spoilage of refrigerated food is very much dependent on temperature (2, 3).

Crystal Violet Tetrazolium Agar Base is used for the detection of gram-negative psychrophilic bacteria causing food spoilage. It is based on the formulation by Olson (4) and recommended by APHA (5) for detecting gram-negative psychrotrophic bacteria.

Casein enzyme hydrolysate and yeast extract provide various nitrogenous nutrients to the organisms while dextrose serves as the carbon source. Crystal violet inhibits most of the gram-positive organisms and therefore inclusion of crystal violet in the medium does not affect the growth of psychrotrophic organisms, which are mostly gram-negative.

Standard methods for the detection of gram-negative psychrotrophic bacteria should be followed (5).

#### Methodology

Suspend 23.5 grams of dehydrated media powder in 1000 ml distilled water. Mix thoroughly & heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 50°C and aseptically add 5 ml of sterile 1% solution of 2, 3, 5-Triphenyl Tetrazolium Chloride (MS 2057). Mix well and pour into sterile Petri plates.

#### Quality Control

##### Appearance

Cream to greyish yellow homogeneous free flowing powder.

##### Gelling

Firm, comparable with 1.5% Agar gel.

##### Colour and Clarity

Light purple coloured, clear to slightly opalescent gel forms in Petri plates.



Dehydrated Culture Media  
Bases / Media Supplements

#### Reaction

Reaction of 2.35% w/v aqueous solution at 25°C. pH : 7.0±0.2

#### pH Range

6.80-7.20

#### Cultural Response

DM 1586: Cultural characteristics observed with added 1% T.T.C. solution (MS 2057) after an incubation at 20-30°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
<b>Cultural Response</b> <i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	good-luxuriant	>=50%	maroon
<i>Staphylococcus aureus</i> ATCC 25923	>=10 <sup>3</sup>	inhibited	0%	
<i>Yersinia enterocolitica</i> ATCC 27729	50-100	good-luxuriant	>=50%	maroon

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Mossel D. A. A., and Zwart H., 1960, J. Appl. Bacteriol., 23:185-188.
2. Elliott R. P. and Michener H. D., 1965, U.S. Dept. Agr. Tech. Bull.No. 1320, p. 110, Washington, D.C.
3. Tomkin R. B., 1973, Food Technol., 27:54.
4. Olson H. C., 1963, J. Dairy Sci., 46:362.
5. Speck M. L., (Ed.), 1984, Compendium of Methods for the Microbiological Examination of Foods, 2nd Ed., APHA, Washington, D.C.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specification for identity and performance parameters.

