

## Technical Information

### Candida Medium

**Product Code: DM 1104**

**Application:** - Candida Medium is used for cultivating *Candida* species.

### Composition\*\*

Ingredients	Gms / Litre
Mycological peptone	2.500
Dextrose	5.000
Disodium hydrogen phosphate	5.000
Sodium sulphite	5.000
Bismuth sulphite indicator	3.000
Agar	15.000
Final pH ( at 25°C)	7.6±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Principle & Interpretation

*Candida* is a genus of yeasts responsible for infections such as oropharyngeal candidiasis, vaginal candidiasis and candidemia.

Candida Medium is used for the selective cultivation and differentiation of *Candida species*. Candida Medium was originally developed by Nickerson (1). It is also used for processing and inoculation of specimens like tissues, skin scraping, nails and hair (2, 3).

Mycological peptone in the medium provides essential nitrogenous nutrients while dextrose acts as carbon source and phosphate maintains buffering action of medium. This medium also contains sodium sulphite, which is reduced by *Candida* species to form sulphide. Bismuth in the medium combines with the sulphide to produce brown to black pigmented colonies and zones of dark precipitate in the medium surrounding the colonies of some species. Bismuth sulphite also acts as an inhibitor of bacterial growth. Selectivity of medium is increased by incorporation of penicillin and streptomycin in the medium, which helps to suppress the growth of many bacteria.

Differentiation of *Candida* is based on the growth patterns and pigmentation of isolated colonies.

### Methodology

Suspend 35.5 grams of dehydrated media powder in 1000 ml distilled water. Mix thoroughly & heat to boiling to dissolve the medium completely. DO NOT AUTOCLAVE. Cool to 50 - 52°C and aseptically add 0.3 units of Penicillin and 25 µg Streptomycin per ml of sterile medium. Mix well before pouring into sterile Petri plates.

### Quality Control

#### Appearance

Cream to yellow homogeneous free flowing powder

#### Gelling

Firm, comparable with 1.5% Agar gel

#### Colour and Clarity

Light yellow coloured, clear to slightly opalescent gel forms in Petri plates

#### Reaction

Reaction of 3.55% w/v aqueous solution at 25°C. pH : 7.6±0.2

#### pH Range

7.40-7.80



**Cultural Response**

DM 1104: Cultural characteristics observed after an incubation at 30°C for 24-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery
<i>Candida albicans</i> ATCC 10231	50-100	good-luxuriant	>=50%
<i>Candida tropicalis</i> ATCC 1369	50-100	good-luxuriant	>=50%
<i>Escherichia coli</i> ATCC 25922	>=10 <sup>3</sup>	inhibited	0%

**Storage and Shelf Life****Dried Media:** Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.**Further Reading**

1. Nickerson, 1953, J. Infect. Dis., 93:43.
2. Haley, Trandel and Coyle, 1980, Cumitech 11, Practical Methods for Culture and Identification Of Fungi In The Clinical Mycology Laboratory, Coord Ed., Sherris, ASM, Washington, D.C.
3. Emmons, Binford, Utz and Kwon-Chung, 1977, Medical Mycology, 3rd Ed., W. B. Saunders Co., Philadelphia.

**Disclaimer :**

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specification for identity and performance parameters.