

## Technical Information

### MUG MFC Agar

#### Product Code: DM 2387

**Application:** - MUG MFC Agar is used for cultivation and enumeration of faecal coliforms by the membrane filter technique at elevated temperature.

#### Composition\*\*

Ingredients	Gms / Litre
Tryptose	10.000
Proteose peptone	5.000
Yeast extract	3.000
Bile salts	1.500
Sodium chloride	5.000
4-Methylumbelliferyl $\beta$ -D-Glucuronide (MUG)	0.100
Agar	15.000
Final pH ( at 25°C)	7.4 $\pm$ 0.2

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

MUG MFC Agar is MFC Agar with added MUG and is used for cultivating and enumerating faecal coliforms by the membrane filtration technique. Faecal coliforms are organisms present in the gastrointestinal tract of warm blooded animals. Presence of faecal coliforms in water is an indication of faecal contamination. MFC Agar, formulated by Geldrich et al (1) is recommended for the detection of faecal coliforms in water samples by the membrane filter technique (2). *Escherichia coli* is a member of the faecal coliform group, which possess the enzyme  $\beta$ -glucuronidase. This enzyme specifically cleaves 4-Methylumbelliferyl- $\beta$  -D-Glucuronide (MUG) to form a fluorescent end product 4-Methylumbelliferone. This end product fluoresces under long wave UV light (366 nm) (1).

Tryptose and proteose peptone supply carbon, nitrogen, vitamins and minerals necessary for the growth of faecal coliforms. Yeast extract provides B-complex vitamins, which stimulate bacterial growth. Bile salts prevent the growth of gram-positive microorganisms. Incorporation of MUG in the medium permits rapid detection of *E. coli*, when the medium is observed for fluorescence under long-wave UV light.

#### Methodology

Suspend 39.6 grams of dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat to boil to dissolve the medium completely. DO NOT AUTOCLAVE. Shake well before pour into sterile Petri plates.

#### Quality Control

##### Appearance

Cream to yellow homogeneous free flowing powder

##### Gelling

Firm, comparable with 1.5% Agar gel

##### Colour and Clarity

Yellow coloured clear to slightly opalescent gel forms in Petri plates

##### Reaction

Reaction of 3.96% w/v aqueous solution at 25°C. pH : 7.4 $\pm$ 0.2

**pH Range**

7.20-7.60

**Cultural Response**

DM2387: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours

Organism	Inoculum (CFU)	Growth	Recovery	Fluorescence (under UV)
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	>=50%	positive
<i>Enterococcus faecalis</i> ATCC 29212	>=10 <sup>3</sup>	inhibited	0%	-
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	luxuriant	>=50%	negative
<i>Shigella flexneri</i> ATCC 12022	50-100	luxuriant	>=50%	negative

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and the prepared medium at 2 -8°C. Use before expiry date on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Geldreich E. E., Clark H. F., Huff C. B. and Best L. C., 1965, J. Amer. Water Works Assoc. 57:208.
2. Eaton A. D., Clesceri L. S. and Greenberg A. E., (Eds.), 1995, Standard Methods for the Examination of Water and Wastewater, 19th Ed., American Public Health Association, Washington, D.C.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performances parameters.