

## Technical Information

### Actidione Agar Base without Actidione

#### Product Code: DM 1058

**Application:** - Actidione Agar Base without Actidione is recommended for the enumeration and detection of bacteria in specimens containing large number of yeasts and moulds.

#### Composition\*\*

Ingredients	Gms / Litre
Casein enzymic hydrolysate	5.000
Yeast extract	4.000
Dextrose	50.000
Monopotassium phosphate	0.550
Potassium chloride	0.425
Calcium chloride	0.125
Magnesium sulphate	0.125
Ferric chloride	0.0025
Manganese sulphate	0.0025
Bromo cresol green	0.022
Agar	15.000
Final pH ( at 25°C)	5.5±0.2

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

Actidione Agar was formulated by Green and Gray (1) which may be recommended for microbiological investigation during brewing and baking. Actidione® (Cycloheximide) at a concentration of 0.001% permits the growth of bacteria and enhance the growth of most yeasts and moulds except dermatophytes. This medium may be used for the estimation of bacterial contamination of pitching yeast. Addition of penicillin or streptomycin may be used for selective isolation of dermatophytes.

Casein enzymic hydrolysate acts as source of nitrogen while yeast extract is a rich reservoir of vitamins. Dextrose in high amount along with mineral salts at acidic pH favours sugar fermentation.

#### Methodology

Suspend 75.25 grams of dehydrated culture media in 1000 ml distilled water. Mix thoroughly & heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45°C and aseptically add 10 mg cycloheximide to the medium. Shake well before pouring into sterile Petriplates.

**Warning:** Cycloheximide (Actidione) is very toxic. Avoid skin contact or aerosol formation and inhalation.

#### Quality Control

##### Appearance

Light yellow to light green homogeneous free flowing powder

##### Gelling

Firm, comparable with 1.5% Agar gel

##### Colour and Clarity

Greenish blue clear to slightly opalescent gel forms in Petri plates

##### Reaction

Reaction of 7.53% w/v aqueous solution at 25°C. pH : 5.5±0.2



Dehydrated Culture Media  
Bases / Media Supplements

#### pH Range

5.30-5.70

#### Cultural Response

DM1058: Cultural characteristics observed after an incubation at 25-30°C for 40-48 hours.

Organism	Inoculum (CFU)	Growth (Plain)	Recovery	Growth (w/ Actidione)	Recovery (w / Actidione)
<b>Cultural Response</b> <i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	>=70%	good-luxuriant	>=50%
<i>Lactobacillus fermentum</i> ATCC 9338	50-100	luxuriant	>=70%	good-luxuriant	>=50%
<i>Proteus mirabilis</i> ATCC 25933	50-100	luxuriant	>=70%	good-luxuriant	>=50%
<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	luxuriant	>=70%	inhibited	0%
<i>Saccharomyces uvarum</i> ATCC 9080	50-100	luxuriant	>=70%	inhibited	0%

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Green, S.R. and Gray, P.P., 1950, Wallerstein Lab. Communication, 13:357.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performs parameters.

