

Technical Information

Glucose Agar

Product Code: DM 2746

Application: - Glucose Agar is used for differentiation of *Enterobacteriaceae* in urine, water and food samples.

Composition**

Ingredients	Gms / Litre
Tryptone	2.000
Glucose	10.000
Sodium chloride	5.000
Yeast extract	1.000
Potassium hydrogen phosphate	0.300
Bromothymol blue	0.080
Agar	2.500
Final pH (at 25°C)	7.1±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Glucose agar is recommended for the differentiation of *Enterobacteriaceae* in urine, water and food. It differentiates species on the basis of glucose fermentation. The *Enterobacteriaceae* are gram- negative chemoautotroph that posses both respiratory and fermentative metabolism.

Tryptone and Yeast extract provide essential nutrients for growth: nitrogen, vitamins, minerals and amino acids; Glucose is the fermentable carbohydrate providing carbon and energy; Sodium chloride maintains the osmotic balance; Bromothymol blue is a pH indicator. Inoculate and incubate at 35 ± 2°C for 18 - 24 hours. The glucose fermenting microorganisms produce yellow colour (acid) and the non-fermenting ones, blue colour or colourless.

Methodology

Suspend 20.88 grams of dehydrated media powder in 1000 ml distilled water. Mix thoroughly & heat to boil to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes and dispense as desired.

Quality Control

Appearance

Light yellow to light green homogeneous free flowing powder

Colour and Clarity

Blue green coloured, clear to slightly opalescent gel forms in tubes as butts.

Reaction

Reaction of 2.08% w/v aqueous solution at 25°C. pH : 7.1±0.2

pH Range

6.90-7.30

Cultural Response

DM 2746: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.



Dehydrated Culture Media
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth	Acid production
Cultural Response			
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	good	Positive reaction(Colour changes to yellow)
<i>Escherichia coli</i> ATCC 25922	50-100	good	Positive reaction(Colour changes to yellow)
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	good	Positive reaction(Colour changes to yellow)

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and prepared media at 2-8°C. Use before expiry date on label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Waisbren, Carr and Dunnett, 1951, Am. J. Clin. Path., 21:884.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performances parameters.

