

Technical Information

MacConkey Agar w/ Magnesium Sulphate

Product Code: DM 2612

Application: - MacConkey Agar w/ Magnesium Sulphate is recommended for isolating and differentiating gram-negative bacilli while suppressing the swarming of most *Proteus* species.

Composition**

Ingredients	Gms / Litre
Pancreatic digest of gelatin	10.000
Yeast extract	10.000
Lactose	10.000
Oxgall	5.000
Magnesium sulphate	0.200
Neutral red	0.075
Agar	12.000
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

MacConkey Agar with Magnesium sulphate is a differential medium recommended for isolation and cultivation of gram-negative enteric organisms and gram-positive cocci from samples suspected of containing these organisms. This medium also limits the swarming of *Proteus* species due to the omission of sodium chloride (3).

MacConkey Agar is the earliest selective and differential medium for cultivation of enteric microorganisms from a variety of clinical specimens (1, 2). The original medium contains protein, bile salts, sodium chloride and two dyes. Selectivity of the medium is attributed to crystal violet and bile salts. MacConkey Agar contains lactose with neutral red to distinguish the lactose-fermenting coliforms from the lactose non-fermenting *Salmonella* and *Shigella* groups (3).

Pancreatic digest of gelatin and yeast extract provide necessary nitrogen sources. Lactose serves as fermentable carbohydrate. Oxgall serves to improve the selectivity of the medium. Gram-negative bacteria usually grow well on the medium and are differentiated by their ability to ferment lactose. Lactose fermenting strains grow as red or pink coloured colonies and may be surrounded by a zone of acid precipitated bile as found in the case of *Escherichia coli*. The red colour is due to production of acid from lactose, absorption of neutral red and a subsequent colour change of the neutral red dye when the pH of medium falls below 6.8. Lactose non-fermenting strains, such as *Shigella* and *Salmonella* appear as colourless and transparent colonies.

Methodology

Suspend 47.27 grams of dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat to boiling with gentle swirling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Avoid overheating. Cool to 45 - 50°C and Shake well before pour into sterile Petri plates.

Quality Control

Appearance

Light yellow to pink homogeneous free flowing powder

Gelling

Firm, comparable with 1.2% agar gel.

Colour and Clarity

Red coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 4.72% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH Range

7.20-7.60

Cultural Response

DM2612: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours (upto 48 hours).

Organism	Inoculum (CFU)	Growth	Recovery	Colour of Colony
<i>Enterococcus faecalis</i> ATCC 29212	50-100	good	40-50%	pink-red
<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant	≥50%	pink - red with bile precipitate
<i>Proteus mirabilis</i> ATCC 12453	50-100	good-luxuriant	≥50%	Colourless, no swarming
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	good-luxuriant	≥50%	colourless
<i>Salmonella Typhi</i> ATCC 6539	50-100	Good	40-50%	colourless
<i>Shigella flexneri</i> ATCC 12022	50-100	good-luxuriant	≥50%	colourless

Storage and Shelf Life

Dried Media: Store below 30°C and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. MacConkey A., 1900, The Lancet, ii.20.
2. MacConkey A., 1905, J. Hyg, 8:333
3. Collee J. G., Fraser A. G., Marmion B. P., Simmons A., (Eds.), Mackie and McCartney, Practical Medical Microbiology, 1996, 14th Edition, Churchill Livingstone

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performances parameters.