

Technical Information

Nutrient Gelatin

Product Code: DM 1060S

Application: - Nutrient Gelatin is used for detection of gelatin liquefaction by proteolytic microorganisms.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	5.000
Meat extract	3.000
Gelatin	120.000
Sodium chloride	30.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Nutrient Gelatin is prepared as per the formulation recommended by BIS (1). Gelatin liquefaction is one of the essential test for the differentiation of enteric bacilli (2). This medium can also be used for the microbial plate counts of water.

Peptic digest of animal tissue and meat extract supply nutrients for the growth of nonfastidious organisms. Organisms produce gelatinase, a proteolytic enzyme active in the liquefaction of gelatin.

To test gelatin liquefaction the strains are stab inoculated in Nutrient Gelatin. Many species require prolonged incubation (3, 4) for gelatin liquefaction. Gelatin is solid at 20°C or less temperature and liquid at 35°C or higher temperature. Gelatin liquefies at about 28°C, so incubation is carried out at 35°C but kept in a refrigerator for about 2 hours before interpretation of the results (3). Liquefaction of gelatin occurs on the surface layer, so care should be taken not to shake the tubes (5). Control is run along with every testing as gelling ability of gelatin varies (3) and also the gelatin concentration should not exceed 12% as it may inhibit growth (6). For plate counts of water, the incubation is carried out at 20-22°C up to 30 days.

Methodology

Suspend 158 grams of dehydrated powder media in 1000 ml of warm (50°C) water. Mix thoroughly & heat to 50°C to dissolve the medium completely. Dispense into test tubes. Sterilize by autoclaving at 15 lbs pressure (121°C) for 12 minutes.

Quality Control

Appearance

Cream to yellow coloured homogeneous free flowing slightly coarse powder

Gelling

Semisolid, comparable with 12% Gelatin gel.

Colour and Clarity

Light amber coloured clear to slightly opalescent gel forms in tubes

Reaction

Reaction of 15.8% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH Range

6.80-7.20

Cultural Response

DM 1060S: Cultural characteristics after 1 to 7 days at 35 - 37°C.

Organism	Inoculum (CFU)	Growth	Gelatinase
<i>Clostridium perfringens</i> ATCC 12924	50-100	good-luxuriant	Positive reaction
<i>Bacillus cereus</i> ATCC 10876	50-100	good-luxuriant	Positive reaction
<i>Bacillus subtilis</i> ATCC 6633	50-100	good-luxuriant	Positive reaction
<i>Escherichia coli</i> ATCC25922	50-100	good-luxuriant	Negative reaction
<i>Proteus vulgaris</i> ATCC 13315	50-100	good-luxuriant	Positive reaction
<i>Staphylococcus aureus</i> ATCC 25923	50-100	good-luxuriant	Positive reaction

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Bureau of Indian Standards IS : 5887 (Part IV) 1976.
2. Ewing, 1986, Edwards and Ewings Identification of Enterobacteriaceae, 4th ed., Elsevier Science Publishing Co., Inc. New York.
3. Cawan S. and Steel K., 1966, Manual for the Identification of Medical Bacteria, Cambridge University Press, Pg. 19, 27-28, 116 and 156.
4. Lautrop H., 1956, Acta Pathol. Microbiol. Scand., 39:357.
5. Frobisher M., 1957, Fundamentals of Microbiology, 6th ed., W.B. Saunders Co., Philadelphia, P:239.
6. Branson D., 1972, Methods in Clinical Bacteriology, Springfield, Ill, pg 21.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications