

Technical Information

Acetate Differential Agar, Modified

Product Code: DM 1339F

Application:- Acetate Differential Agar, Modified is recommended for the differentiation of *Shigella* species from *Escherichia coli* in accordance with FDA BAM, 1998.

Composition**

Ingredients	Gms / Litre
Sodium acetate	2.000
Sodium chloride	5.000
Magnesium sulphate	0.200
Ammonium phosphate	1.000
Dipotassium phosphate	1.000
Bromothymol blue	0.080
Agar	20.000
Final pH (at 25°C)	6.70±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Acetate Differential Agar, Modified (DM 1339F) is recommended for the differentiation of *Shigella* species from *E. coli* in accordance with FDA BAM, 1998 (1). Shigellosis, although commonly regarded as waterborne, is also a food borne disease majorly caused to humans. It is spread among humans by food handlers with lesser personal hygiene. *Escherichia coli* is widely distributed in the intestine of humans and is an important facultative anaerobe present in the colon area of a healthy individual.

Acetate Differential Agar, Modified was formulated by Trabulsi and Ewing (2), by modifying Citrate Medium of Simmons (3). Most of the bacteria can utilise citrate and acetate as the carbon sources for growth in the presence of organic nitrogen, not in the absence of it. This difference in growth is helpful in differentiating *Shigella* from other closely related organisms such as *E. coli* (4). *E. coli* grows well within 24-48 hours in this media indicated by formation of blue colour ($5 < (> , <) > 6$). Magnesium sulphate is an essential ion. Sodium chloride maintains osmotic equilibrium and phosphates maintain the pH.

Aseptically weigh 25 g sample into 225 ml *Shigella* Broth Base (DM 2326) supplemented with novobiocin. Incubate jars under anaerobic conditions at 44.0°C in a water bath for 20 hrs. This can further be streaked on to a MacConkey agar plate (DM 1081D). Incubate for 20 h at 35°C. After Gram staining, the culture can be proceeded for biochemical confirmation. Inoculate the cultures into slants of Acetate Differential Agar, Modified and incubate overnight at 35°C. Acetate utilization is indicated by formation of blue colour, which is due to the utilization of sodium acetate and subsequent formation of an alkaline reaction detected by the presence of bromothymol blue indicator. *Shigella* is negative and do not show blue colouration, where as *E. coli* is positive (1).

Methodology

Suspend 29.28 grams of dehydrated media in 1000 ml distilled water. Mix thoroughly heat to boiling to dissolve the medium completely. Distribute in tubes in sufficient amounts to give butt and slant. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Allow the tubes to cool in a slanted position.

Quality Control

Appearance

Cream to light green homogeneous free flowing powder.

Gelling

Firm, comparable with 2.0% agar gel.

Colour and Clarity

Emerald green coloured clear to slightly opalescent gel forms in tubes as slants.

Reaction

Reaction of 2.92% w/v aqueous solution at 25°C. pH : 6.7±0.2

pH Range

6.50-6.90

Cultural Response

DM 1339F: Cultural characteristics observed after an incubation at 25-30°C for up to 1-7 days.

Organism	Inoculum (CFU)	Growth	Acetate utilization
Cultural Response			
<i>Citrobacter freundii</i> ATCC 8090	50-100	good-luxuriant	positive reaction, blue colour
<i>Enterobacter cloacae</i> ATCC23355	50-100	good-luxuriant	positive reaction, blue colour
<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant	positive reaction, blue colour
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	good-luxuriant	positive reaction, blue colour
<i>Proteus vulgaris</i> ATCC 13315	>=10 ³	inhibited	-
<i>Salmonella Arizonae</i> ATCC 13314	50-100	good-luxuriant	positive reaction, blue colour
<i>Salmonella Typhi</i> ATCC 19430	50-100	poor	negative reaction green colour
<i>Shigella sonnei</i> ATCC 25931	50-100	none-poor	negative reaction, no change, medium remains green

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Prepared Media: 2-8°C in sealable plastic bags for 2-5 days.

Further Reading

1. FDA, U.S. 1998. Bacteriological Analytical Manual. 8 ed. Gaithersburg, MD: AOAC International.
2. Trabulsi. and Ewing. 1962. Public Health Lab, 20.
3. Simmons. 1926. J. Infect. Dis, 39.
4. Cordaro, J.T. and Ball, R.J. 1966. Applied Microbiology, 14(6): 886-887.
5. Ewing. 1986. Edwards and Ewings Identification of Enterobacteriaceae. 4 ed. N.Y: Elsevier Science Pub. Co., Inc.
6. Talukder, K. A., Islam, M. A., Dutta, D.K., Hasan, F., Sada, A., Nair, G. and Bnd Sack, D. A. 2002. J. Clin. Microbiol, 40.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specification for identity and performance parameters.