

Technical Information

Oxytetra Glucose Yeast Agar Base (OGYE Agar Base)

Product Code: DM 1639

Application: - Oxytetra Glucose Yeast Agar Base (OGYE Agar Base) is used for isolation and enumeration of yeasts and / or moulds from foods.

Composition**

-		
Ingredients	Gms / Litre	
Yeast extract	5.000	
Dextrose (Glucose)	20.000	
Agar	12.000	
Final pH (at 25°C)	7.0±0.2	
1		

^{**}Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

OGYE Agar Base is formulated by Mossel et al for the selective isolation and enumeration of yeast and moulds from foods (1, 2). They found that addition of Oxytetra selective supplement to a neutral pH medium increased the recovery / count of yeast and moulds as compared to acidified medium.

Acidic media are not completely suitable for counting yeasts and moulds in foods since yeast cells, stressed by heat do not tolerate the acidic conditions necessary to prevent the bacterial contamination. Yeast and mould growth is often limited by the presence of acid-tolerant bacterial flora. Therefore it is evident that more active media and different selective agents are needed in order to deal with various kinds of foodstuffs, incubation conditions and types of microorganisms to be studied. Under certain conditions and when testing certain foods like milk and milk products, the use of oxytetracycline alone was not sufficient to obtain reliable yeast and mould counts.

Yeast extract supplies essential growth nutrients. Dextrose acts as carbon and energy source. Low pH helps to reduce the bacterial flora. Oxytetracycline makes the medium more selective by inhibiting the growth of Lactobacilli encountered in milk and milk-products at low pH. The choice of a suitable media for enumeration of yeasts and moulds greatly depends on the nature of foodstuffs to be tested and the organisms that grow on them. These media remain bacteriostatic when inoculated with not greater than 1 ml of a 10-1 food dilution and incubation at 22°C. The number of yeasts or moulds is calculated per one gram or 1 ml of sample under investigation by multiplying the number of colonies with the dilution factor. Lactic acid bacteria are inhibited on this medium.

Type of specimen

Food samples

Specimen Collection and Handling:

For food samples, follow appropriate techniques for sample collection and processing as per guidelines (3).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions:

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. Due to nutritional variation, some strains of fungi may show poor growth.





Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the recommended temperature.

Methodology

Suspend 18.5 grams of dehydrated powder media in 500 ml distilled water. Mix thoroughly & heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add reconstituted contents of one vial of Oxytetra Selective Supplement (MS1032). Shake well before pour into sterile Petri plates.

Quality Control

Appearance

Cream to light yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.2% Agar gel.

Colour and Clarity

Light amber coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 3.7% w/v aqueous solution at 25°C. pH: 7.0±0.2

pH Range

6.80-7.20

Cultural Response

DM1639: Cultural characteristics observed with added 1 vial of Oxytetra Selective Supplement (MS1032), after an incubation at 25-30°C after 2-5 days.

Organism	Inoculum (CFU)	Growth	Recovery
Cultural Response # Aspergillus brasiliensis ATCC 16404 (00053*)	50-100	good-luxuriant	-
Candida albicans ATCC 10231 (00054*)	50-100	good-luxuriant	>=50%
Escherichia coli ATCC 25922 (00013*)	>=10³	inhibited	0%
Saccharomyces cerevisiae ATCC 9763 (00058*)	50-100	good-luxuriant	>=50%
Saccharomyces uvarum ATCC 9080	50-100	good-luxuriant	>=50%

Key: (#) Formerly known as Aspergillus niger (*) Corresponding WDCM numbers.

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle inorder to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (4, 5).





Further Reading

- 1.Mossel D. A. A., Kleynen-Semmeling H. M., Vincentie H., Beerens H. and Catsaras M., 1970, J. Appl. Bacteriol., 33:454
- 2. Mossel D. A. A., Visser M. and Mengerink W. H. J., 1962, Lab. Pract. 11:109.
- 3. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
- 4. Isenberg, H.D. Clinical Microbiology Procedures Handb0ook. 2nd Edition.
- 5. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Disclaimer:

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performens parameters.

