

Technical Information

Broth Medium R (Lactose monohydrate sulphite Medium)

Product Code: DM 2287B

Application: - Lactose monohydrate sulphite Medium is used for detection and enumeration of *Clostridium* in pharmaceutical products as per British Pharmacopoeia, 2009.

Composition**

Ingredients	Gms / Litre
Pancreatic digest of casein	5.000
Yeast extract	2.500
Sodium chloride	2.500
Lactose monohydrate	10.000
Cystine hydrochloride	0.300
Final pH (at 25°C)	7.1±0.1

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Lactose Monohydrate Sulphite Medium base is prepared as per the British Pharmacopoeia (1) and is cited as Medium R. This medium is useful in semi-quantitative test for presence of *Clostridium perfringens* in pharmaceutical products where the level of this species is a criterion of quality.

The medium contains pancreatic digest of casein and yeast extract, which supplies essential nitrogenous compounds for *Clostridia*. Lactose monohydrate acts as carbon or fermentable carbohydrate source. Cystine hydrochloride rich in sulphur content provides reduced conditions. Sodium metabisulphite and ferric ammonium citrate act as indicators of sulphite reduction, indicated by blackening of the medium.

Methodology

Suspend 19.8 grams of dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat if necessary to dissolve in tubes containing inverted Durham's tubes. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and add filter-sterilized solution of 1.2% sodium metabisulphite R (0.5ml) and 1.0% ferric ammonium citrate (0.5ml) to each tube.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity

Light amber coloured clear solution without any precipitate

pH Range

7.00-7.20

Growth Promotion Test

Growth Promotion is carried out in accordance with British pharmacopoeia

Cultural Response

DM2287B: Cultural characteristics after an incubation at 46±0.5°C for 24-48 hours.



Dehydrated Culture Media
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth	H ₂ S	Gas
<i>Clostridium perfringens</i> ATCC 13124	50-100	luxuriant	Positive reaction, blackening of medium	Positive reaction
<i>Clostridium sporogenes</i> ATCC 19404	50-100	luxuriant	negative reaction no blackening of medium	Positive reaction

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. British Pharmacopoeia, 2009, The Stationery office British Pharmacopoeia.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performances parameters.

