

Technical Information

Listeria Oxford Medium Base

Product Code: DM 2145

Application: - Listeria Oxford Medium with supplements is recommended for isolation of *Listeria* species from pathological specimen.

Composition**

Ingredients	Gms / Litre
Peptone, special	23.000
Lithium chloride	15.000
Sodium chloride	5.000
Corn starch	1.000
Esculin	1.000
Ammonium ferric citrate	0.500
Agar	10.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Listeria monocytogenes is the only species of the genus *Listeria* that is important as a human pathogen. *Listeria seeligeri*, *Listeria welshimeri* and *Listeria ivanovii* have been related with animal diseases. In any case, all the species are pathogenic between the ovine and bovine cattle. Positive diagnosis of listeriosis can be obtained only by the isolation and cultivation of the responsible bacteria from blood or CSF samples of the affected organisms. Listeria Oxford Medium Base is based on the formulation described by Curtis et al (1) for isolation of *L. monocytogenes* from clinical and food specimens.

Peptone special serves as the source of essential nutrients to the organisms. Corn starch serves to neutralize the toxic metabolites formed. Lithium chloride and the antibiotics inhibit gram-negative bacteria and most gram-positive organisms but certain strains of Staphylococci may grow as esculin negative colonies. Cycloheximide is used to reduce fungal contamination; cefotetan and phosphomycin are inhibitors of bacterial overgrowth. Acriflavin, colistin sulphate and lithium chloride inhibit bacteria other than *Listeria* species. Alternatively moxalactam (MS1126) can be added which inhibits both gram-positive and gram-negative bacteria. *L. monocytogenes* hydrolyzes esculin to esculetin and dextrose. Esculetin reacts with ferric ions and produces black zones around the colonies. Although the selectivity of the medium is enough to allow the isolation and differentiation by direct surface inoculation, a previous dilution of the inoculum is advisable or even more when the sample is highly polluted.

The techniques for isolation vary with the material under examination (2). For all specimens selective and cold enrichment is recommended (3, 4). For faecal and biological specimens, the sample is homogenized in 0.1% Peptone Water (DM1028) and 0.1 ml amount is either directly plated on Listeria Selective Medium or inoculated into the Selective Enrichment Broth and incubated at 30°C for 7 days and then further inoculated on Listeria Selective Medium. For food and environmental samples selective enrichment is generally used.

For isolation of Listeria from food (milk and milk products), add 25 ml or 25 grams of sample to 225 ml of Listeria Enrichment Broth, UVM (DM1890A). Homogenize and mix carefully. Incubate for 48 hours at 30°C. Streak the enriched cultures onto Listeria Oxford medium Base and incubate aerobically for 48 hours at 37°C. Take 5 typical colonies (esculin positive) and inoculate onto Soyabean Casein Digest Medium (DM1290). Incubate for 24 hours and then use these colonies for biochemical confirmation.

Methodology

Suspend 27.75 grams of dehydrated powder media in 500 ml distilled water. Mix thoroughly & heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add the rehydrated contents of 1 vial of Oxford Listeria Supplement (MS1071) or 1 vial of Listeria Moxalactam Supplement (MS2126). Mix well before pouring into sterile Petri plates.

Warning: Lithium chloride is harmful. Avoid bodily contact and inhalation of vapours. On contact with skin, wash with plenty of water immediately.

Quality Control

Appearance

Light yellow to dark yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.0% Agar gel.

Colour and Clarity

Dark amber coloured clear to slightly opalescent gel with a blue cast forms in Petri plates

Reaction

Reaction of 5.55% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH Range

6.80-7.20

Cultural Response

DM2145: Cultural characteristics observed with added Oxford Listeria Supplement (MS2071) or Listeria Moxalactam supplement (MS2126), after an incubation at 35-37°C for 24-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Esculin Hydrolysis
<i>Bacillus subtilis</i> ATCC 6633	≥10 ³	inhibited	0%	-
<i>Enterococcus faecalis</i> ATCC 29212	≥10 ³	inhibited	0%	-
<i>Enterococcus hirae</i> ATCC 10541	≥10 ³	inhibited	0%	-
<i>Escherichia coli</i> ATCC 25922	≥10 ³	inhibited	0%	-
<i>Listeria monocytogenes</i> ATCC 19111	50-100	luxuriant	≥50%	positive reaction, blackening of medium around the colony
<i>Listeria monocytogenes</i> ATCC 19112	50-100	luxuriant	≥50%	positive reaction, blackening of medium around the colony
<i>Listeria monocytogenes</i> ATCC 19117	50-100	luxuriant	≥50%	positive reaction, blackening of medium around the colony
<i>Staphylococcus aureus</i> ATCC 25923	50-100	luxuriant	40-50%	negative reaction

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and prepared medium at 2-8°C. Use before expiry period on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.



Dehydrated Culture Media
Bases / Media Supplements

Further Reading

1. Curtis G. D. W., Mitchell R. G, King A. F., Griffin E. J., 1989, Lett. Appl. Microbiol., 8:95
2. Van Netten P., Peroles I., Van de Mosdik A., Curtis G. D. W., Mossel D. A. A, 1988, Int. J. Food Microbiol., 6:187.
3. Hayes P. S, Feeley J. L, Groves L. M, Ajello G. W. and Fleming D. W, 1986, Appl. Environ. Microbiol., 51:438.
4. Fernandez G. J. F., Dominguez R. L., Vazzuez B. J. A., Rodriguez F.E. F., Briones D. V., Blanco L. J. L., Suarez F. G., 1986, Can. J. Microbiol., 32:149.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performances parameters.

