

Technical Information

Salmonella Differential Agar (Twin Pack)(RajHans Medium)

Product Code: DM 2078

Application: - Salmonella Differential Agar is recommended for identification and differentiation of *Salmonella* species from members of *Enterobacteriaceae*, especially *Proteus* species.

Composition**

Ingredients	Gms / Litre
Part A	-
Peptone, special	8.000
Yeast extract	2.000
Sodium deoxycholate	1.000
B. C. Indicator	2.000
Agar	12.000
Part B	-
Propylene glycol	10.000
Final pH (at 25°C)	7.3±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Salmonella Differential Agar is slight modification of original formulation of Rambach (1) used for differentiation of *Salmonella* species from *Proteus* species and other enteric bacteria. Production of acid from propylene glycol is a novel characteristic of *Salmonella* species and is utilized in these media. Many of the media such as SS Agar, XLD Agar recommended for the identification and differentiation of *Salmonella* species (2) are based on lactose fermentation and hydrogen sulphide production.

Peptone special and yeast extract supports the luxuriant growth of bacteria while sodium deoxycholate inhibits gram-positive organisms rendering the medium selective for enteric microorganisms. The BC indicator turns pink in presence of acid produced from propylene glycol. Lactose fermenting ability is determined by using an indicator, which can detect the presence of enzyme β -galactosidase. Lactose fermenting (β -galactosidase producing) bacteria yield blue violet coloured colony (3). *Salmonellae* produce acid from propylene glycol and on combining with the pH indicator gives typical pink red colonies. Other enteric gram-negative bacteria form colourless colonies. *Salmonella* Typhimurium and *Salmonella* Enteritidis produce pink to red colonies. Specimen should be enriched in an appropriate selective enrichment broth. This enriched culture is then inoculated on Salmonella Differential Agar and incubated at 35-37°C for 24-48 hours.

Methodology

Suspend 10 grams of dehydrated media fluid of **Part B** in 1000 ml distilled water. Add 25 grams of dehydrated powder media of **Part A**. Mix well and heat to boil to dissolve the medium completely. DO NOT AUTOCLAVE. Cool to 45-50°C. Mix well before pouring into sterile Petri plates.

Quality Control

Appearance

Part A: Light yellow to light pink homogeneous free flowing powder

Part B: Colourless viscous solution

Gelling

Firm, comparable with 1.2% Agar gel.

Colour and Clarity

Light orange coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 2.5% w/v aqueous solution of **Part A** at 25°C. pH : 7.3±0.2

pH Range

7.10-7.50

Cultural Response

DM 2078: Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
Cultural Response				
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	>=50%	blue-green
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	luxuriant	>=50%	blue-violet
<i>Proteus mirabilis</i> ATCC 25933	50-100	luxuriant	>=50%	colourless
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	luxuriant	>=50%	pink-red
<i>Salmonella Enteritidis</i> ATCC 13076	50-100	luxuriant	>=50%	pink-red
<i>Salmonella Typhi</i> ATCC 6539	50-100	luxuriant	>=50%	colourless
<i>Shigella flexneri</i> ATCC 12022	50-100	luxuriant	>=50%	colourless
<i>Staphylococcus aureus</i> ATCC 25923	>=10 ³	inhibited	0%	-

Storage and Shelf Life

Dried Media: Store dehydrated powder and prepared medium at 2-8°C. Use before expiry period on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Rambach A., 1990, Appl Environ. Microbiol., 56:301.
2. Eaton A. D., Clesceri L. S., Rice E. W. and Greenberg A W., (Eds.), 2005, Standard Methods for the Examination of Water and Wastewater, 21st Ed., APHA, Washington, D.C.
3. Greenwald R., Henderson R.W. and Yappaw S., 1991, J. Clin. Microbiol. 29:2354.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.