

Technical Information

R-3A Broth

Product Code: DM 2688

Application: - R-3A Broth is used for sub culturing microorganisms from potable water.

Composition**

Ingredients	Gms / Litre
Casein acid hydrolysate	1.000
Yeast extract	1.000
Biopeptone	1.000
Dextrose	1.000
Starch soluble	1.000
Dipotassium phosphate	0.600
Magnesium sulphate anhydrous	0.048
Sodium pyruvate	0.600
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

R-3A Agar is a subculture medium which is used to subculture the organisms recovered on nutritionally deficient R-2A Agar (1). R-3A Broth is similar in composition to R-3A Agar, except agar.

The total bacterial count of drinking water is determined by plate count on a nutritionally rich medium. However all organisms present are not able to grow on them, either because they are slow growers or because they can grow on that media (1). For this reason a nutritionally reduced medium was described. R-2A Agar is a modification of this medium (2,3).

Many bacteria from natural waters, which contain limited nutrients at ambient temperature, grow best on the media with less nutrient levels. They grow better at the temperatures below the routine laboratory incubation temperatures of 35 to 37°C (3).

These media contain casein acid hydrolysate, yeast extract, biopeptone as source of essential growth factors required for metabolism of the bacteria. Dextrose act as the energy source. Starch acts as a neutralizer that neutralizes any toxic metabolites, if present. Phosphate buffers the medium while sodium pyruvate supplies additional nutrition. Magnesium sulphate serves as a source of ions. Due to the presence of the above mentioned ingredients these media allow the growth of stressed and chlorine tolerant bacteria present in treated waters.

Methodology

Suspend 6.25 grams of dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat if necessary to dissolve the medium completely. Dispense into tubes. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. DO NOT OVERHEAT.

Quality Control

Appearance

Cream to yellow coloured homogeneous free flowing powder.

Colour and Clarity

Yellow coloured clear solution in tubes.

Reaction

Reaction of 0.625% w/v aqueous solution at 25°C. pH : 7.2±0.2

pH Range

7.00-7.40

Cultural Response

DM 2688: Cultural characteristics observed *by using standard ATCC cultures after an incubation at 35-37°C for 24-72 hours.

Organism	Inoculum (CFU)	Growth
<i>Candida albicans</i> ATCC 10231	50-100	good-luxuriant
<i>Enterococcus faecalis</i> ATCC 29212	50-100	good-luxuriant
<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant
<i>Salmonella Enteritidis</i> ATCC 13076	50-100	good-luxuriant
<i>Salmonella Typhi</i> ATCC 6539	50-100	good-luxuriant

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Reasoner and Geldreich, 1985, Appl. Environ. Microbiol., 49:1.
2. Stark and McCoy. 1938. Zentralbl. Bacteriol. Parasitenkd. Infektionskr. Hyg. Abt.2 98 : 201
3. Collins and Willoughby, 1962, Arch. Microbiol., 43:294.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specification for identity and performance parameters.