

Technical Information

Nutrient Agar w/ Tyrosine

Product Code: DM 1561F

Application: - Nutrient Agar w/ Tyrosine is recommended for cultivation and enumeration of *Bacillus cereus* in water and food in accordance with FDA BAM, 1998.

Composition**

Ingredients	Gms / Litre
Beef extract	3.000
Peptone	5.000
Agar	15.000
Tyrosine	5.000
Final pH (at 25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Nutrient Agar w/ Tyrosine is recommended for cultivation and enumeration of *Bacillus cereus* in water and food in accordance with FDA BAM, 1998(1) *Bacillus cereus* is an aerobic spore-forming bacterium that is commonly found in soil, on vegetables, and in many raw and processed foods. *B. cereus* food poisoning may occur when foods are prepared and held without adequate refrigeration for several hours before serving, with *B. cereus*. Foods incriminated in past outbreaks include cooked meat and vegetables, boiled or fried rice, vanilla sauce, custards, soups, and raw vegetable sprouts. The organism can be identified by its ability to hydrolyze tyrosine in the medium. Peptone and beef extract supply essential nutrients for the growth of the organism. Agar acts as the solidifying agent. Tyrosine acts as a source of amino acid which is hydrolyzed by *Bacillus* species.

Prepare 1:10 dilutions of 50 g of the sample in Butterfield's phosphate-buffered dilution water. Plate count of *B.cereus* can be done on MYP (DM1636F) agar plates from appropriate dilutions. *B.cereus* gives pink coloured colonies on MYP agar. Suspected colonies are subcultured into Nutrient agar (DM1561F). Inoculate entire surface of tyrosine agar slant with mm loopful of culture from Nutrient agar. Incubate slants 48 h at 35°C. Positive results are indicated by the zone of clearance in and around the bacterial growth, indicating hydrolysis. Examine negative slants for obvious signs of growth, and incubate for a total of 7 days before considering as negative (1). This media is used in the confirmation of other species of *Bacillus* such as *B. cereus*, *B. thuringiensis*, *B. mycoides*, *B. weihenstephanensis*, *B. anthracis* and *B. megaterium* and also for *Streptomyces* and *Nocardia* species (2, 3).

Methodology

Suspend 28.00 grams of dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Shake well and dispense 3.5 ml into sterile tubes with frequent mixing. Keep in slanted position and cool rapidly to prevent separation of tyrosine.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel



Dehydrated Culture Media
Bases / Media Supplements

Colour and Clarity

Yellow coloured clear to slightly opalescent gel forms in slants(may shows some white particles after solidification)

Reaction

Reaction of 2.8% w/v aqueous solution at 25°C. pH : 6.8±0.2

pH Range

6.60-7.00

Cultural Response

DM1561F: Cultural characteristics observed after an incubation at 35-37°C for 48 hours up to 7days.

Organism	Growth	Tyrosine hydrolysis
<i>Bacillus cereus</i> ATCC 10876	good-luxuriant	positive reaction, clearing of medium in and around the bacterial growth
<i>Bacillus thuringiensis</i> ATCC 1079	good-luxuriant	positive reaction, clearing of medium in and around the bacterial growth
<i>Escherichia coli</i> ATCC 25922	good	negative reaction, no clear zones

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2 -8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. FDA, U.S. 1998. Bacteriological Analytical Manual. 8 ed. Gaithersburg, MD: AOAC International.
2. L arone. 1995. Medically important fungi: a guide to identification. 3 ed. Washington, D.C: ASM.
3. Murray, P. R., Baron, E. J., Jorgensen, J. H., Pfaller, M. A. and Tenover, R. C. 2003. Manual of Clinical Microbiology. 8 ed. Washington, D.C: ASM.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.

