

Technical Information

Coagulase Mannitol Agar Base

Product Code: DM 1272

Application: - Coagulase Mannitol Agar Base is recommended for the primary isolation and differentiation of pathogenic Staphylococci from clinical specimens or for classifying pure cultures.

Composition**

Ingredients	Gms / Litre
Brain heart infusion	5.000
Casein enzymic hydrolysate	10.500
Papaic digest of soyabean meal	3.500
Sodium chloride	3.500
Mannitol	10.000
Bromo cresol purple	0.020
Agar	14.500
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

The genus *Staphylococcus* comprises 28 accepted or proposed species, 14 of which may be encountered in human clinical specimens. Staphylococci are generally found on the skin and mucous membranes of humans and other animals. Some of the pathogenic staphylococci in both humans and animals produce an enzyme called coagulase and detection of this enzyme is used in the laboratory to identify these organisms (6).

This media are used for the isolation of *Staphylococcus aureus* from clinical specimens and for differentiation of *S.aureus* from other species on the basis of coagulase production and mannitol fermentation. Chapman for the first time introduced a medium for selective isolation and differentiation of Staphylococci (1). Tellurite-glycine media were formulated by Zebovitz et al (2) and Marwin (3) for selectively isolating coagulase-positive Staphylococcal species. Present medium is based on Esber and Faulconer formulation (5). Mutant or old cultures of *S.aureus* may be weak coagulase producers. They should be freshly sub cultured and rechecked. *Escherichia coli* ferments mannitol and may be weakly coagulase positive. Coagulase production is dependent on the presence of a fermentable sugar like mannitol in this case. It is also dependent on the presence of a protein factor in the brain heart infusion and blood plasma (4). When mannitol is fermented, the pH of the medium surrounding the coagulase positive colonies drops. This drop in pH is indicated by the change in colour of the bromocresol purple indicator, which turns yellow and exhibits yellow zones around the colonies.

An opaque area of coagulated plasma forms around the colonies of coagulase positive organisms. *Staphylococcus epidermidis* is coagulase negative and mannitol non-fermenting species, which does not change the colour of the medium. Coagulase negative species may ferment mannitol and produce a yellow zone around the colonies but an opaque zone will not be formed.

Methodology

Suspend 47.02 of grams dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 118 - 121°C (12-15 lbs pressure respectively) for 15 minutes. Cool to 45 - 50°C. Add 7 - 15% v/v sterile, pre-tested, rabbit plasma to basal medium. Mix well and pour into sterile Petri plates.

Quality Control

Appearance

Light yellow to light grey homogeneous free flowing powder.

Gelling

Firm, comparable with 1.45% Agar gel.

Colour and Clarity

Purple coloured, slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 4.7% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH Range

7.20-7.60

Cultural Response

DM 1272: Cultural characteristics observed with added 7-15% v/v sterile pretested, rabbit plasma at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Mannitol fermentation	Coagulase production
<i>Staphylococcus aureus</i> ATCC 25923	50-100	luxuriant	≥70%	positive reaction, yellow colour	positive reaction, colonies surrounded by opaque negative reaction, no opaque zone
<i>Staphylococcus epidermidis</i> ATCC 12228	50-100	luxuriant	≥70%	negative reaction, purple colour	negative reaction, no opaque zone

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Chapman, 1944, J. Bacteriol., 48:113
2. Zebovitz, Evans and Nivens, 1955, J. Bacteriol., 70:686.
3. Marwin, 1958, Am. J. Clin. Pathol., 30:470.
4. Schaub and Merrit, 1960, Bull. Johns Hopkins Hosp., 106:25.
5. Esber and Faulconer, 1959, Am. J. Clin. Pathol., 32:192.
6. Koneman E. W., Allen S. D., Janda W. M., Schreckenberger P. C., Winn W. C. Jr., 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4th Ed., J. B. Lippincott Company

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specification for identity and performance parameters.