

## Technical Information

### Clostridium Brazier Agar Base

**Product Code: DM 2803**

**Application:** - Clostridium Brazier Agar Base is recommended as a selective medium for the isolation and differentiation of *Clostridium difficile* with added supplements.

### Composition\*\*

Ingredients	Gms / Litre
Peptone special	23.000
Sodium chloride	5.000
Starch, soluble	1.000
Sodium bicarbonate	0.400
Dextrose	1.000
Sodium pyruvate	1.000
Cysteine HCL	0.500
Haemin	0.010
Vitamin K	0.001
L-Arginine	1.000
Sodium pyrophosphate	0.250
Sodium succinate	0.500
Cholic acid	1.000
p-Hydroxyphenylacetic acid	1.000
Agar	12.000
Final pH ( at 25°C)	7.0±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Principle & Interpretation

The spectrum of disease caused by *Clostridium difficile* (a pathogenic *Clostridium* affecting the bowel) ranges from pseudomembranous colitis (PMC) through antibiotic associated colitis (AAC). It also includes chronic inflammatory bowel diseases, post-operative diarrhoea and non-antibiotic associated diarrhoea (1). Smith and King (2) first reported the presence of *C. difficile* in human infections.

Clostridium Brazier Agar Base was developed by Jon Brazier (3) based on similar work carried out by Ken Phillips and Paul Levett. Many pathological labs including Anaerobe Reference Unit are using this medium for isolating *C. difficile*. Typical characteristics of *C. difficile* appears on this medium after 24 hours on anaerobic incubation at 35-37°C. *C. difficile* appears as grey, opaque, flat raised colonies generally circular but may tend to elongate, which on further incubation upto 48 hours may result in lighter grey or may impart white centre to the medium and form opaque colonies, 4-6 mm in diameter. Typical Gram stain morphology of *C. difficile* may not be seen in colonies taken from this medium due to the presence of antibiotics. The selective agents in Clostridium difficile supplement (MS1010), D-cycloserine and cefoxitin used in this medium inhibits the growth of the majority of *Enterobacteriaceae* and also *Enterococcus faecalis*, Staphylococci, gram negative anaerobic bacilli and *Clostridium* species other than *C. difficile* which may be found abundantly in samples. The Egg Yolk Emulsion (MS 2045) added to the medium helps to differentiate *C. difficile* from lecithinase positive Clostridia. Addition of lysed horse blood to the base enhances recognition of colony fluorescence when cultures are examined using UV light. Cholic acid present in the medium promotes spore germination following shock treatment, and p-hydroxyphenylacetic acid to enhance production of p-cresol, a distinctive metabolite of *C. difficile*.

## Methodology

Suspend 47.66 grams of dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat to boil to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Aseptically add rehydrated contents of 2 vials of Clostridium Difficile Supplement (MS 2010), 40 ml of Egg Yolk Emulsion (MS 2045) together with 10 ml lysed horse blood. Mix well and pour into sterile Petri plates.

## Quality Control

### Appearance

Cream to yellow homogeneous free flowing powder

### Gelling

Firm, comparable with 1.2% Agar gel

### Colour and Clarity

**Basal medium:** Light amber coloured clear to slightly opalescent gel. After addition of Egg yolk emulsion (MS 2045) and 10 ml lysed horse blood: Tan coloured opaque gel forms in Petri plates.

### Reaction

Reaction of 4.76% w/v aqueous solution at 25°C. pH : 7.0±0.2

### pH Range

6.80-7.20

### Cultural Response

DM2803: Cultural characteristics observed under anaerobic condition with added Clostridium Difficile Supplement (MS 2010), Egg yolk Emulsion (MS 2045) and 10 ml of lysed horse blood, after an incubation at 35-37°C for 48 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony	Lecithinase activity
<b>Cultural Response</b> <i>Clostridium difficile</i> ATCC 11204	50-100	good-luxuriant	≥50%	greyish-white, opaque flat colonies	negative
<i>Escherichia coli</i> ATCC 25922	≥10 <sup>3</sup>	inhibited	0%	-	-

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Collee J. G., Fraser A. G., Marmion B. P., Simmons A., (Eds.), Mackie and McCartney, Practical Medical Microbiology, 14th Ed., Churchill Livingstone.
2. Smith L. D. S. and King E. O., 1962, J. Bacteriol., 84:65.
3. Brazier J S (1993) Role of the Laboratory in Investigations of Clostridium difficile Diarrhoea. Clinical Infectious Diseases 16 (4) S228-33.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performances parameters.