

Technical Information

MUG Sorbitol Agar

Product Code: DM 2205

Application: - MUG Sorbitol Agar is recommended for the isolation and identification of enteropathogenic *Escherichia coli* associated with infant diarrhea by fluorogenic method.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	17.000
Proteose peptone	3.000
D-Sorbitol	10.000
Bile salts mixture	1.500
Sodium chloride	5.000
Neutral red	0.030
Crystal violet	0.001
4-Methylumbelliferyl β -D-Glucuronide (MUG)	0.100
Agar	13.500
Final pH (at 25°C)	7.1 \pm 0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Escherichia coli serotype O157:H7 is a human pathogen associated with hemorrhagic colitis. Most organisms of the faecal flora ferment sorbitol and appear pink on this medium. MUG Sorbitol Agar is a modification of MacConkey Agar using sorbitol instead of lactose. MUG Sorbitol Agar is used for detecting or differentiating enteropathogenic *E. coli* (EPEC) in water by a fluorogenic method. The distinction of EPEC from other groups of pathogenic *E. coli* isolated from patients' stools involves serological and cell culture assays. EPEC causes watery diarrhea and bloody diarrhea. Watery diarrhea is associated with attachment and physical alteration of the integrity of the intestine. Bloody diarrhea is associated with attachment of acute tissue destructive process mediated by a toxin called shiga toxin or verotoxin. Shiga toxin is cell associated rather than excreted. Hence the detection or differentiation of this organism is vital from public health point of view.

Among the other strains of *E. coli*, the enteropathogenic strain lacks the sorbitol degrading ability within 48 hours of incubation. Moreover it does not synthesize the enzyme glucuronidase and hence there is no fluorescence production by this strain when MUG is present in the medium (1). Bile salts mixture and crystal violet in the medium prevent most of the gram- positive organisms, which accompany the specimen many times. Sorbitol, a polyhydric alcohol corresponding to glucose, serves as a substrate to determine the cleavage of sorbitol by sorbitol degrading microorganisms. Sorbitol degrading microorganisms produce pink to red colonies while sorbitol negative colonies are colourless. MUG (4-Methyl-umbellifery β -D-Glucuronide) is converted into a fluorescent product 4-Methyl-umbelliferone by the β -D-glucuronidase-producing organisms. However enteropathogenic *E. coli* (in contrast to commensal *E. coli* strains) does not synthesize this enzyme and thus when its colonies are exposed to long wave UV light, no fluorescence is observed. The plates are exposed to ammonia fumes to increase fluorescence as suggested by Freir and Hartman (4).

It has reported that some *Enterobacteriaceae* and *Pseudomonas aeruginosa* are inhibited on this medium when incubated in a CO₂-enriched atmosphere (2). The colour of sorbitol- positive colonies can fade, making them hard to distinguish from sorbitol-negative colonies (3).

Methodology

Suspend 50.13 grams of dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Shake well before pour into sterile Petri plates.

Quality Control

Appearance

Light yellow to pink homogeneous free flowing powder

Gelling

Firm, comparable with 1.35% Agar gel.

Colour and Clarity

Purplish red coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 5.01% w/v aqueous solution at 25°C. pH : 7.1±0.2

pH Range

6.90-7.30

Cultural Response

DM 2205: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Colour of colony	Sorbitol	Fluorescence (under UV)*
<i>Escherichia coli O157:H7</i>	50-100	good-luxuriant	colourless	negative	negative
<i>Escherichia coli ATCC25922</i>	50-100	good-luxuriant	pink-red	positive	positive
<i>Staphylococcus aureus ATCC 25923</i>	>=10 ³	inhibited	-	-	-

Key: * Fluorescence can be visualized by addition of NaOH solution or exposure to ammonia fumes.

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Szabo R. A., Todd E. C. and Jean A., 1986, J. Food Prot., 10:768.
2. Mazura- Reetz, Neblett G. T. and Galperin J. M., 1979, Abstr. C 179, p. 339, Abst. Annu. Med. Am. Soc., Microbiol.
3. Adams, 1991, Clin. Lab. Sci., 4 :19
4. Freir T.A. and Hartman P.A. (1987) Appl. Env. Microbiol. 53. 1246-1250

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.