

Technical Information

Liver Broth

Product Code: DM 1928

Application: - Liver Broth is recommended for examination of food for saccharolytic, putrefactive, mesophilic and thermophilic anaerobes.

Composition**

Ingredients	Gms / Litre
Liver, infusion from	23.000
Peptic digest of animal tissue	10.000
Liver tissues (extracted)	30.000
Dipotassium phosphate	1.000
Final pH (at 25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Scar used Liver Broth for the examination of Canners sugar for hydrogen swells caused by thermophilic anaerobes (1) and also for maintaining pure cultures of aerobes and anaerobes.

This medium contains liver particles which support luxuriant growth for saccharolytic or putrefactive mesophilic and thermophilic anaerobes. A 20% w/v solution of the sugar steamed for 30 minutes is inoculated into Liver Broth, sealed with agar. The standard proposed was a maximum of 1 positive tube in six tubes, with 20 ml inocula and incubated for 72 hours at 56°C. Some organisms like *Clostridium thermosaccharolyticum* produces gas which often pushes the agar plug towards the top of the tube and some organisms digest the solid liver tissues. The medium should be used on the same day of preparation as the stored medium may absorb the air and then re-steaming is necessary which darkens the medium.

Methodology

Suspend 64 grams of dehydrated powder media in 1000 ml distilled water. Soak for 15 minutes with occasional stirring. Dispense in 18 mm diameter tubes to a depth of 50 mm so that bottom of the tube is filled with liver tissues. Sterilize by autoclaving at 10 lbs pressure (115°C) for 20 minutes. Cool, inoculate and seal with a layer of sterile 2% agar solution.

Quality Control

Appearance

Brown coloured granules

Colour and Clarity

Medium amber coloured, clear to slightly opalescent supernatant over insoluble granules

Reaction

Reaction of 6.4% w/v aqueous suspension at 25°C. pH : 6.8±0.2

pH Range

6.60-7.00

Cultural Response

DM1928: Cultural characteristics observed after an incubation at 55-57°C for 48-72 hours.



Dehydrated Culture Media
Bases / Media Supplements

Cultural Response

Organism	Inoculum (CFU)	Growth
Cultural Response <i>Cl.thermosaccharolyticum</i> ATCC 7956	50-100	luxuriant

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use freshly prepared medium. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Scarr M. P., 1958, DSIR, Proc. 2nd Internat. Symp. Food Microbiol., 1957, HMSO London, pp-29.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performances parameters.

