

Technical Information

Eugonic LT 100 Medium Base w/o Tween 80

Product Code: DM 2513

Application: - Eugonic LT 100 Medium Base is used for the cultivation of fastidious microorganisms like *Haemophilus*, *Neisseria*, *Pasturella*, *Brucella* and *Lactobacillus* species.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	15.000
Papaic digest of soyabean meal	5.000
Glucose	5.500
Sodium chloride	4.000
Sodium sulphite	0.200
L-Cystine	0.700
Egg lecithin	1.000
Triton X-100	1.000
Agar	15.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Eugonic LT 100 Medium Base was developed by Pelczar and Vera (1) for cultivation of fastidious organisms like *Brucella*. Eugon media were developed to obtain eugonic (luxuriant) growth of fastidious microorganisms like *Brucella* which are otherwise difficult to cultivate (3). The unenriched medium supports rapid growth of lactobacilli associated with cured meat products, dairy products and other foods. Eugonic media is quite similar to Tryptone Soya Agar (DM 1290) but more bacterial propagation is expected on Eugonic media. Organisms like *Bordetella* and *Neisseria* form minute colonies on Tryptone Soya Agar (DM 1290). They may become large on Eugon Media because large amount of sulfur and carbon sources have been added in addition to the Tryptone Soya Agar (DM 1290) formula. Eugonic LT 100 Medium w/o Tween 80 can be used for growth of a variety of fastidious microorganisms like *Neisseria*, *Francisella* and *Brucella*.

Casein enzymic hydrolysate and papaic digest of soyabean meal provide the nitrogen, vitamins and amino acids, which supports the growth of fastidious microbial species. The high concentration of glucose is the energy source for rapid growth of bacteria. L-Cystine and sodium sulphite are added to stimulate growth. Sodium chloride maintains the osmotic balance of the media. The high carbohydrate content along with high sulfur (cystine) content improves growth with chromogenicity (2). Lecithin and polysorbate 80 in Eugonic LT 100 Medium w/o Tween 80 neutralize antimicrobial agents hence this medium can be used as a neutralizing diluent.

Methodology

Suspend 47.40 grams of dehydrated media in 1000 ml distilled water containing 5 grams of polysorbate 80 (Tween 80). Mix thoroughly & heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Shake well before pour into sterile Petri plates.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity

Yellow coloured, clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 4.74% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH Range

6.80-7.20

Cultural Response

DM 2513: Cultural characteristics observed with added 5-10% sterile defibrinated blood after an incubation at 35-37°C for 48 hours (fungal cultures incubated at 25-30°C).

Organism	Inoculum (CFU)	Growth	Recovery
<i>Bacillus pumilus</i> ATCC 14884	50-100	good (with 0.1% starch)	50-70%
<i>Brucella abortus</i> ATCC 4315	50-100	good (under 3-5% CO ₂)	50-70%
<i>Candida albicans</i> ATCC 26790	50-100	good	50-70%
<i>Lactobacillus fermentum</i> ATCC 9338	50-100	good	50-70%
<i>Neisseria meningitidis</i> ATCC 13090	50-100	good	50-70%
<i>Streptococcus pneumoniae</i> ATCC 6303	50-100	good- luxuriant(under 3-5% CO ₂)	>=70%
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	good- luxuriant(under 3-5% CO ₂)	50-70%
<i>Staphylococcus aureus</i> ATCC 25923	50-100	good-luxuriant	>=70%
<i>Staphylococcus aureus</i> ATCC 6538	50-100	good-luxuriant	>=70%
<i>Candida albicans</i> ATCC 10231	50-100	good	50-70%
<i>Bacillus subtilis</i> ATCC 6633	50-100	good	50-70%
<i>Pseudomonas aeruginosa</i> ATCC 9027	50-100	good	50-70%
<i>Escherichia coli</i> ATCC 8739	50-100	good-luxuriant	50-70%

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared media at 2-8°C. Use before expiry period on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Pelczar and Vera J., 1949, Milk Plant Monthly 38:30
2. Frank H. A., 1955, J. Bacteriol., 70:269.
3. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams & Wilkins, Baltimore, Md.



Dehydrated Culture Media
Bases / Media Supplements

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specification for identity and performance parameters.

