

Technical Information

Rinsing Fluid

Product Code: DM 2622

Application: - Rinsing Fluid is recommended as a rinsing fluid in the membrane filtration procedure.

Composition**

Ingredients	Gms / Litre
Peptone from meat	5.000
Meat extract	3.000
Polysorbate 80	1.000
Final pH (at 25°C)	6.9±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Rinsing fluid largely complies with the formulation prescribed in the recommendations of the United States Pharmacopoeia (1). After filtration it is often necessary to rinse the membrane filter in order to remove residues of liquid sample materials. If the sample contains higher hydrocarbons such as vaseline, paraffin, etc. or fats, the use of rinse fluid is recommended.

This fluid contains balanced concentrations of nutrients, which prevent the microorganisms, retained by the filter, from being exposed to physiological shock, thus being capable to grow further rapidly. The detergent polysorbate 80 ensures emulsification of carbohydrates and fats without seriously affecting the microorganisms. If the sample contains large quantities of these compounds, additional up to 9.0 g/litre of polysorbate 80 (Tween 80) can be added in accordance with the USP recommendations before the broth is filtered. After filtering the liquid sample, rinse the filter 3 times with 100 ml portions of the membrane-filter rinse fluid, and then complete the test in the usual way.

Methodology

Suspend 9 grams of dehydrated powder media in 1000 ml distilled water. If desired together with up to 9 g/litre of polysorbate 80 (Tween 80) to i.e , filter until clear. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Quality Control

Appearance

Cream to yellow coloured homogeneous free flowing powder.

Colour and Clarity

Yellow coloured clear solution without any precipitate.

Reaction

Reaction of 0.9%w/v aqueous solution at 25°C pH : 6.9±0.2

pH Range

6.70-7.10

Cultural Response

DM2622: Cultural characteristics observed after an incubation at 35 -37°C for 18 - 24 hours. Recovery is carried out on Standard Nutrient Agar No.1 (DM2210)



Dehydrated Culture Media
Bases / Media Supplements

Organism	Growth
<i>Staphylococcus aureus</i> ATCC 25923	Good
<i>Streptococcus pyogenes</i> ATCC 12344	Good
<i>Enterococcus faecalis</i> ATCC 29212	Good
<i>Citrobacter freundii</i> ATCC 8090	Good
<i>Pseudomonas aeruginosa</i> ATCC 27853	Good
<i>Escherichia coli</i> ATCC 25922	Good

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. The United States Pharmacopoeia, 2006, USP29/NF24. The United States Pharmacopeial Convention. Rockville, MD.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performances parameters.

