

Technical Information

Cryptococcus Differential Agar

Product Code: DM 2814

Application: - Cryptococcus Differential Agar is recommended for a differentiation of *Cryptococcus* species.

Composition**

Ingredients	Gms / Litre
Glucose	20.000
Glycine	0.500
DL- Tryptophan	2.000
Potassium dihydrogen phosphate	4.000
Magnesium sulphate	2.500
Thiamine HCl	0.005
Trypan Blue	0.030
Agar	15.000
Final pH (at 25°C)	5.4±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Cryptococcus is the etiological agent of cryptococcosis, a systemic mycosis of humans and animals with a worldwide distribution.

Cryptococcosis (earlier called European blastomycosis) commonly starts following inhalation of the organism, which is considered opportunistic infections as it affects mainly immunosuppressed individuals. (3)

Cryptococcus Differential Agar was based on the formulation of m-FDTG medium except the sugar fructose was replaced by glucose as it supported better growth of *Cryptococcus* species. Glucose supports growth as well as strong pigment production by nearly all *C. gattii* strains. *C. gattii* can while *C. neoformans* cannot assimilate D-tryptophan (1), thereby producing a brown diffusible pigment (4). Pigmentation is not apparent on the first day of growth but is usually noticeable after 5 days of incubation, intensity gradually increases with time after 2-3 weeks. (2).

Glycine serves as a sole source of carbon and nitrogen which is utilized by *Cryptococcus gattii* *Cryptococcus laurentii* and not by *Cryptococcus neoformans*. Salts in the medium help in pigment induction by D-tryptophan. Pigment production was more intense at 25-30°C as compared to 37°C. Dyes in media for the isolation of fungi have not been commonly utilized, although many such media are available for the isolation of bacteria. Trypan blue medium allows suspected *C. neoformans* colonies to be subcultured before mold overgrowth becomes a problem (5).

Methodology

Suspend 44.04 grams of dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat to boiling to dissolve the medium completely. DO NOT AUTOCLAVE. Cool to 50°C and pour into sterile Petri plates.

Quality Control

Appearance

Light yellow to yellow with bluish tinge homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel



Dehydrated Culture Media
Bases / Media Supplements

Colour and Clarity

Light blue coloured, opalescent gel with white precipitate forms in Petri plates

Reaction

Reaction of 4.4 % w/v aqueous solution at 25°C. pH : 5.4±0.2

pH Range

5.20-5.60

Cultural Response

DM2814: Cultural characteristics observed after an incubation at 25- 30°C for 5 to 6 days.

Organism	Inoculum (CFU)	Growth	Colony Characteristics
----------	----------------	--------	------------------------

Cultural Response

<i>Cryptococcus neoformans</i> ATCC 32045	50-100	luxuriant	Light blue, dry colony
---	--------	-----------	------------------------

<i>Cryptococcus laurentii</i> ATCC 18803	50-100	luxuriant	Brown, dry colony
--	--------	-----------	-------------------

<i>Cryptococcus gattii</i> ATCC MYA- 4566	50-100	luxuriant	Brown, mucoid colony
---	--------	-----------	----------------------

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Baro, T., J.m. Torres-Rodriguez, M.H. De Mendoza, Y. Morera, and C.Alia. 1998 First identification of autochthonous *Cryptococcus neoformans* var. *gattii* isolated from goats with predominantly severe pulmonary disease in Spain. *J.Clin. Microbiol.* 36:458-461
2. Chaskes,S., Frases, S., Cammer, M., Gerfen, G, and Casadevall, A. (2008).Growth and Pigment Production on D-Tryptophan Medium by *Cryptococcus gattii* , *Cryptococcus neoformans* , and *Candida albicans* . *J. Clin. Microbiol.* 46 : 255-264.
3. Misral, V.C, and Randhawa. H.S (2000). Occurrence and Significance of *Cryptococcus neoformans* in Vegetables and Fruits. *The Indian Journal of Chest Diseases 6 Allied Sciences.* 42:317-322.
4. Mukamurangwa,P., C.Raes- Wuytack, and C. De Vroey.1995. *Cryptococcus neoformans* var. *gattii* can be separated from ! var. *neoformans* by its ability to assimilate D-tryptophan. *J.Med. Vet. Mycol.*33:419–420.
5. Racicot,T.A, and Bulmer,G.S. (1985). Comparison of Media for the Isolation of *Cryptococcus neoformans*. *Appl. And Environ. Microbiol.* 50(2): 548-549.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performances parameters.

