

Technical Information

Sabouraud Cycloheximide Chloramphenicol Agar

Product Code: DM 1664

Application: - Sabouraud Cycloheximide Chloramphenicol Agar is recommended for selective isolation and cultivation of fungi.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	10.000
Dextrose	20.000
Chloramphenicol	0.040
Cycloheximide	0.500
Agar	15.000
Final pH (at 25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Sabouraud Dextrose Agar was originally formulated by Sabouraud (1) and further modified by Emmons (2) by reducing dextrose content and adjusting the pH close to neutral.

Peptic digest of animal tissue is the source of nitrogenous growth factors while dextrose supplies an energy source for the growth of microorganisms. The media can be rendered selective for fungi by antibiotics such as Chloramphenicol (4) and Cycloheximide (5), which inhibit some bacteria as well as some saprophytic and pathogenic fungi. This medium inhibits fungi like *Cryptococcus neoformans*, *Aspergillus*, *Nocardia*, certain *Candida* species but allow the dermatophytes to grow well.

Methodology

Suspend 45.54 grams of dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat to boil to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Shake well and pour into sterile Petri plates.

Caution: Cycloheximide is very toxic. Avoid skin contact or aerosol formation and inhalation.

Some pathogenic fungi may produce infective spores, which are easily dispersed in air, so examination should be carried out in safety cabinet.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder.

Gelling

Firm, comparable with 1.5% Agar gel.

Colour and Clarity

Light amber coloured clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 4.5% w/v aqueous solution at 25°C. pH : 6.8±0.2

pH Range

6.60-7.00



Dehydrated Culture Media
Bases / Media Supplements

Cultural Response

DM 1664: Cultural characteristics observed after an incubation at 25-30°C for 2-3 weeks.

Organism	Inoculum (CFU)	Growth	Recovery
* <i>Aspergillus brasiliensis</i> ATCC 16404	50-100	none-poor	-
<i>Candida albicans</i> ATCC 10231	50-100	poor-fair	<=20%
<i>Escherichia coli</i> ATCC 25922	>=10 ³	inhibited	0%
<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	none-poor	<=20%
<i>Trichophyton mentagrophytes</i> ATCC 9533	50-100	luxuriant	-
<i>Trichophyton rubrum</i> ATCC 28191	50-100	luxuriant	-

*Key: Formerly known as *Aspergillus niger*

Storage and Shelf Life

Dried Media: Store dehydrated medium and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Sabouraud R., 1892, Ann. Dermatol. Syphilol., 3:1061.
2. Emmons C., Binford C., Uty J. and Kwon-Chung, 1970, Medical Mycology, 2nd ed., Philadelphia: Lea and Febiger.
3. Diagnostic Procedures, 1963, 4th ed., APHA
- 4 Ajello L., 1957, J. Chron. Dis., 5:545. ,,
- 5 MacFaddin J. F., 1985, Media For Isolation-Cultivation Identification - Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore. ,,

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specificatons for identity and performens parameters.

