

Technical Information

Fluconazole Testing Medium (Twin Pack)

Product Code: DM 2209

Application: - Fluconazole Testing Medium is used for fluconazole susceptibility testing using *Candida* species.

Composition**

Ingredients	Gms / Litre
Part A	-
Agar	10.000
Part B	-
Dextrose	19.980
Potassium dihydrogen phosphate	1.990
Ammonium sulphate	4.990
L-Glutamine	0.580
Magnesium sulphate anhydrous	0.990
Sodium chloride	0.200
Calcium chloride	0.200
L-Lysine monohydrochloride	0.073
Valine	0.047
L-Arginine monohydrochloride	0.042
L-Histidine	0.023
DL-Methionine	0.0189
Tryptophan	0.020
Nicotinic acid	0.00079
Inositol	0.00397
Pyridoxine hydrochloride	0.00079
Boric acid	0.00099
Calcium D-pantothenic acid	0.00079
Aneurine hydrochloride	0.00079
Manganous sulphate	0.00079
Zinc sulphate	0.0014
p-Amino benzoic acid (PABA)	0.000395
Riboflavin	0.000395
Ferric chloride	0.000395
Cupric sulphate	0.00012
Biotin crystalline	0.000004
Folic acid	0.000395
L-Isoleucine	0.052
Sodium molybdate	0.00047
Potassium iodide	0.0002
L-Leucine	0.052
Threonine	0.0476

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

To carry out standardization of antifungal drug susceptibility assays as a standard reference method, CLSI had recommended broth macro dilution testing of yeasts. Fluconazole Testing Medium is a chemically defined medium specifically developed for the in-vitro testing of fluconazole by using *Candida* species. Inhibitory concentration values obtained by using this medium correlate well with the clinical outcome (1, 2, 3).

The medium contains dextrose and a variety of amino acids, salts and vitamins to support the growth of *Candida* and other fungi.

The inoculum size varies with different fungi. *Candida* species are grown in Sabouraud Dextrose Broth (DM1033) at 37°C for 16-18 hours and then diluted with normal saline to give following dilutions:

Candida albicans ,,,,,, 105 / ml

Candida tropicalis ,,,,, 105 / ml

Candida krusei ,,,,,, 105 / ml

Candida guilliermondii ,,,, 106 / ml

Candida parapsilosis ,,,, 106 / ml

Candida pseudotropicalis 106 / ml

Surface inoculate the above diluted cultures and incubate at 28°C for 48 hours to determine MIC value of fluconazole.

Dermatophytes are grown in Sabouraud Dextrose Agar (DM1063) at 28°C for 5-10 days. The mycelial growth is homogenized in 2 ml of 0.85% saline using glass beads. The density of the suspension is adjusted with 0.85% saline to get a 65% light transmission. Inoculate the plates and incubate for about 5-6 days at 28°C.

Check the control plates to ensure that all isolates have grown adequately and determine the Minimum Inhibitory Concentration (MIC).

Methodology

Part A and **Part B** are sterilized separately as follows:

Part A: - Suspend 2.0 grams of **Part A** in 100 ml distilled water, add 0.1 ml phosphate buffer to adjust the pH to 7.5. Heat to boiling to dissolve the agar particles completely and then sterilize by autoclaving at 115°C for 10 minutes.

Part B: - Suspend 29.31 grams of **Part B** in 900 ml distilled water. Shake well, add 2 gram of sodium bicarbonate, after stirring make up the total volume to 1 litre with distilled water. Sterilize by filtration. The medium can be kept at 4°C for two weeks. Complete medium is prepared by aseptically adding equal volume of molten **Part A** (previously cooled to 50°C) and **Part B**. Mix thoroughly and dispense.

Quality Control

Appearance

Part A: Cream to light yellow homogeneous coarse powder

Part B: White to light yellow homogeneous free flowing powder

Colour and Clarity

Light yellow coloured, opalescent solution may be with fine precipitate

Cultural Response

Cultural characteristics observed after an incubation at 28-30°C for 48 hours .

Cultural Response

Organism	MIC of Fluconazole
----------	--------------------

Cultural Response

<i>Candida albicans</i> ATCC 10231	1.56 µg/ml
------------------------------------	------------



Dehydrated Culture Media
Bases / Media Supplements

Storage and Shelf Life

Dried Media: Store powder medium and prepared medium at 2-8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Hoeprich P. D. and Finn. P. D., 1972, J. Infect, Dis., 126: 353
2. Cook R. A., McIntyre K. A. and Galgiani J. N., 1990, Antimicrob. Agents and Chemother., 34:1542.
3. Pfaller M. A. et al, 1992, Antimicrob. Agents and Chemother., 36:1805.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.

