

Technical Information

Loeffler Serum Medium Base

Product Code: DM 2189

Application: - Loeffler Serum Medium Base with added bovine serum is recommended for the cultivation of *Corynebacterium diphtheriae*.

Composition**

Ingredients	Gms / Litre
Heart muscle, infusion from	0.720
Peptic digest of animal tissue	0.710
Sodium chloride	0.360
Dextrose	0.710
Egg powder	7.500
Final pH (at 25°C)	7.6±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Corynebacterium diphtheriae, also called as Klebs-Loeffler bacillus, is a gram-positive, non-encapsulated, non-sporulated, non-motile facultative anaerobe. It causes infection in humans, leading to diseased condition called diphtheria characterized by an inflammatory lesion and membranous exudates on the mucosa of the upper respiratory tract. *C. diphtheriae* may show abundant volutin in films from a moist Loeffler serum slope. Preliminary culture on Loeffler Agar is required to induce the characteristic production of abundant granules in *C. diphtheriae*. Loeffler Medium was originally devised by Loeffler (1) and was further modified by Perry and Petran (2) and Buck (3). Loeffler medium enhances primary and secondary isolation and cultivation of fastidious pathogenic microorganisms especially from nose and throat. It also restores virulence and other identifying properties (microscopic and colonial) after they have been lost due to prolonged incubation or repeated subculturing. It is also used for demonstration of pigmentation and ascospores.

The high serum content helps in determining proteolytic activity of organisms. Peptic digest of animal tissue, heart muscle infusion, and bovine serum supply the amino acids and other complex nitrogenous substances to support growth of *Corynebacterium*. Dextrose acts as source of fermentable carbohydrate and energy. Sodium chloride helps to maintain the osmotic balance.

Rub the swabs directly over the surface of medium. Following incubation prepare smears from the surface of slope. For testing proteolysis, inoculate slant and prior to incubation, flood the slant with Brewer Thioglycollate Medium (DM 1019). Incubation should be carried out for 3-4 days or much longer for appearance of proteolysis. Loeffler Medium should be used in parallel with Serum Tellurite Agar for selective isolation of *Corynebacterium* (4). Examine cultures and smears stained with Loefflers methylene blue after incubation. Observe for typical cellular morphology of *Corynebacterium* species and for the presence of metachromatic granules which take up the methylene blue dye. Subculture colonies that are catalase-positive and exhibit typical morphology on blood agar, for identification procedures. Observe for pigmentation of specific organisms; e.g., *Pseudomonas aeruginosa* (green) and *Staphylococcus aureus* (yellow to gold). Proteolytic activity is evidenced by destruction of the integrity of the coagulated medium.

Although the production of metachromatic granules on this medium is characteristic of members of the *Corynebacterium* genus, other organisms, such as *Propionibacterium*, some Actinomyces and pleomorphic streptococcal strains display stained granules resembling those of the *Corynebacterium* (4).

Methodology

Suspend 10 grams of dehydrated powder media in 250 ml distilled water at a temperature of 45°C. Mix thoroughly. Add 750 ml of sterile Bovine serum, Shake well and dispense into tubes. Coagulate and sterilize by inspissation for 15 minutes at 80 to 90°C or steaming at 100°C for 10-15 minutes.

Quality Control

Appearance

Light yellow to brownish yellow homogeneous

Colour and Clarity

Basal medium: Light amber coloured clear solution After addition of horse serum and coagulation: Off-white coloured opalescent slants forms in tubes

Reaction

Reaction of basal medium (10gms in 250ml distilled water) at 25°C. pH : 7.6±0.2

pH Range

7.40-7.80

Cultural Response

DM 2189: Cultural characteristics observed with added sterile Bovine serum after an incubation at 35-37°C for 24-48 hours.

Organism

Growth

Cultural Response

Corynebacterium diphtheriae ATCC 11913

fair-good

Corynebacterium diphtheriae type *mitis* *Corynebacterium*

good-luxuriant

diphtheriae type *gravis*

good-luxuriant

Pseudomonas aeruginosa ATCC 10145

good (green colonies with proteolysis)

Staphylococcus aureus ATCC 25923

good (yellow to gold colonies)

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Loeffler, 1887, Zentralbl. Bakteriologie, Parasitenkd., 2:105.
2. Perry and Petran, 1939, J. Lab. Clin. Med., 25:71.
3. Buck, 1949, J. Lab. Clin. Med., 34:582.
4. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.